

HTR1F / 5-HT1F Receptor Antibody (Cytoplasmic Domain)
Rabbit Polyclonal Antibody
Catalog # ALS10188**Specification**

HTR1F / 5-HT1F Receptor Antibody (Cytoplasmic Domain) - Product Information

Application	IHC
Primary Accession	P30939
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42kDa KDa

HTR1F / 5-HT1F Receptor Antibody (Cytoplasmic Domain) - Additional Information**Gene ID** 3355**Other Names**

5-hydroxytryptamine receptor 1F, 5-HT-1F, 5-HT1F, Serotonin receptor 1F, HTR1F, HTR1EL

Target/Specificity

Human 5HT1F Receptor. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage

Long term: -70°C; Short term: +4°C

Precautions

HTR1F / 5-HT1F Receptor Antibody (Cytoplasmic Domain) is for research use only and not for use in diagnostic or therapeutic procedures.

HTR1F / 5-HT1F Receptor Antibody (Cytoplasmic Domain) - Protein Information**Name** HTR1F**Synonyms** HTR1EL**Function**

G-protein coupled receptor for 5-hydroxytryptamine (serotonin). Also functions as a receptor for various alkaloids and psychoactive substances. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate cyclase. Signaling inhibits adenylate cyclase activity.

Cellular Location

Cell membrane; Multi-pass membrane protein

Volume


50 µl

HTR1F / 5-HT1F Receptor Antibody (Cytoplasmic Domain) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HTR1F / 5-HT1F Receptor Antibody (Cytoplasmic Domain) - Images



Anti-5HT1F Receptor antibody ALS10188 IHC of human brain, cortex.

HTR1F / 5-HT1F Receptor Antibody (Cytoplasmic Domain) - Background

G-protein coupled receptor for 5-hydroxytryptamine (serotonin). Also functions as a receptor for various alkaloids and psychoactive substances. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate cyclase. Signaling inhibits adenylate cyclase activity.

HTR1F / 5-HT1F Receptor Antibody (Cytoplasmic Domain) - References

Lovenberg T.W., et al. Proc. Natl. Acad. Sci. U.S.A. 90:2184-2188(1993).
Adham N., et al. Proc. Natl. Acad. Sci. U.S.A. 90:408-412(1993).
Puhl H.L. III, et al. Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases.
Muzny D.M., et al. Nature 440:1194-1198(2006).
Nichols D.E., et al. Chem. Rev. 108:1614-1641(2008).