

**GPR34 Antibody (C-Terminus)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS10317****Specification**

---

**GPR34 Antibody (C-Terminus) - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | IHC                    |
| Primary Accession | <a href="#">Q9UPC5</a> |
| Reactivity        | Human                  |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Calculated MW     | 44kDa KDa              |

**GPR34 Antibody (C-Terminus) - Additional Information****Gene ID** 2857**Other Names**

Probable G-protein coupled receptor 34, GPR34

**Target/Specificity**

Human GPR34. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except GPR82 (100%).

**Reconstitution & Storage**

Long term: -70°C; Short term: +4°C

**Precautions**

GPR34 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

**GPR34 Antibody (C-Terminus) - Protein Information****Name** GPR34**Function**

G-protein-coupled receptor of lysophosphatidylserine (LysoPS) that plays different roles in immune response (PubMed:<a href="http://www.uniprot.org/citations/16460680" target="\_blank">16460680</a>). Acts a damage-sensing receptor that triggers tissue repair upon recognition of dying neutrophils (By similarity). Mechanistically, apoptotic neutrophils release lysophosphatidylserine that are recognized by type 3 innate lymphoid cells (ILC3s) via GPR34, which activates downstream PI3K-AKT and RAS-ERK signaling pathways leading to STAT3 activation and IL-22 production (By similarity). Plays an important role in microglial function, controlling morphology and phagocytosis (By similarity).

**Cellular Location**

Cell membrane; Multi-pass membrane protein

**Tissue Location**

Broadly expressed. Highly expressed on mast cells (PubMed:16460680).


**Volume**

50 µl


**GPR34 Antibody (C-Terminus) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**GPR34 Antibody (C-Terminus) - Images**


Anti-GPR34 antibody IHC of human Soft Tissue, Liposarcoma.



Anti-GPR34 antibody IHC of human Soft Tissue, Liposarcoma.



Anti-GPR34 antibody ALS10317 IHC of human brain, neurons and glia.



Anti-GPR34 antibody IHC of human Lymph Node, Hodgkins Lymphoma.

#### **GPR34 Antibody (C-Terminus) - Background**

Orphan receptor.

#### **GPR34 Antibody (C-Terminus) - References**

Schoneberg T.,et al.Biochim. Biophys. Acta 1446:57-70(1999).  
Marchese A.,et al.Genomics 56:12-21(1999).  
Jacobi F.K.,et al.Hum. Genet. 107:89-91(2000).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).