

## NR2E1 / TLX Antibody (N-Terminus)

Rabbit Polyclonal Antibody Catalog # ALS10552

### **Specification**

## NR2E1 / TLX Antibody (N-Terminus) - Product Information

Application IHC
Primary Accession Q9Y466

Reactivity Human, Rabbit, Monkey, Pig, Horse,

**Bovine**, Dog

Host Rabbit
Clonality Polyclonal
Calculated MW 43kDa KDa

## NR2E1 / TLX Antibody (N-Terminus) - Additional Information

### **Gene ID 7101**

#### **Other Names**

Nuclear receptor subfamily 2 group E member 1, Nuclear receptor TLX, Protein tailless homolog, TII, hTII, NR2E1, TLX

# **Target/Specificity**

Human NR2E1. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

### **Reconstitution & Storage**

Long term: -70°C; Short term: +4°C

#### **Precautions**

NR2E1 / TLX Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

# NR2E1 / TLX Antibody (N-Terminus) - Protein Information

### Name NR2E1

### **Synonyms TLX**

### **Function**

Orphan receptor that binds DNA as a monomer to hormone response elements (HRE) containing an extended core motif half-site sequence 5'-AAGGTCA-3' in which the 5' flanking nucleotides participate in determining receptor specificity (By similarity). May be required to pattern anterior brain differentiation. Involved in the regulation of retinal development and essential for vision. During retinogenesis, regulates PTEN-Cyclin D expression via binding to the promoter region of PTEN and suppressing its activity (By similarity). May be involved in retinoic acid receptor (RAR) regulation in retinal cells.



### **Cellular Location**

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00407}.

### **Tissue Location**


Brain specific. Present in all brain sections tested, highest levels in the caudate nucleus and hippocampus, weakest levels in the thalamus.

# NR2E1 / TLX Antibody (N-Terminus) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# NR2E1 / TLX Antibody (N-Terminus) - Images




Anti-NR2E1 antibody ALS10552 IHC of human brain, neurons and glia.



Anti-NR2E1 / TLX antibody IHC of human Lung, Small Cell Carcinoma.

NR2E1 / TLX Antibody (N-Terminus) - Background





Tel: 858.875.1900 Fax: 858.875.1999

Orphan receptor that binds DNA as a monomer to hormone response elements (HRE) containing an extended core motif half- site sequence 5'-AAGGTCA-3' in which the 5' flanking nucleotides participate in determining receptor specificity (By similarity). May be required to pattern anterior brain differentiation. Involved in the regulation of retinal development and essential for vision. During retinogenesis, regulates PTEN-Cyclin D expression via binding to the promoter region of PTEN and suppressing its activity (By similarity). May be involved in retinoic acic receptor (RAR) regulation in retinal cells.

## NR2E1 / TLX Antibody (N-Terminus) - References

Jackson A., et al. Genomics 50:34-43(1998). Kobayashi M., et al. Mol. Cell. Biol. 20:8731-8739(2000). Hua F., et al. Submitted (AUG-2001) to the EMBL/GenBank/DDBJ databases. Ota T., et al. Nat. Genet. 36:40-45(2004). Mungall A.J., et al. Nature 425:805-811(2003).