10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

GPR44 / CRTH2 Antibody (Extracellular Domain)

Rabbit Polyclonal Antibody
Catalog # ALS10674

Specification

GPR44 / CRTH2 Antibody (Extracellular Domain) - Product Information

Application	**ICC**
Primary Accession	Q9Y5Y4
Reactivity	**Human, Monkey**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**43kDa KDa**

GPR44 / CRTH2 Antibody (Extracellular Domain) - Additional Information

Gene ID 11251

Other Names
Prostaglandin D2 receptor 2, Chemoattractant receptor-homologous molecule expressed on Th2 cells, G-protein coupled receptor 44, CD294, PTGDR2, CRTH2, DL1R, GPR44

Target/Specificity
Human GPR44 / CRTH2. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage
Long term: -70°C; Short term: +4°C

Precautions
GPR44 / CRTH2 Antibody (Extracellular Domain) is for research use only and not for use in diagnostic or therapeutic procedures.

GPR44 / CRTH2 Antibody (Extracellular Domain) - Protein Information

Name PTGDR2

Synonyms CRTH2, DL1R, GPR44

Function
Receptor for prostaglandin D2 (PGD2). Coupled to the G(i)- protein. Receptor activation may result in pertussis toxin-sensitive decreases in cAMP levels and Ca(2+) mobilization. PI3K signaling is also implicated in mediating PTGDR2 effects. PGD2 induced receptor internalization. CRTH2 internalization can be regulated by diverse kinases such as, PKC, PKA, GRK2, GPRK5/GRK5 and GRK6. Receptor activation is responsible, at least in part, in immune regulation and allergic/inflammation responses.

Cellular Location
Cell membrane; Multi-pass membrane protein. Note=Internalized receptors colocalized with

RAB11A.

Tissue Location
Widespread expression. High expression in stomach, small intestine, heart and thymus. Intermediate expression in colon, spinal cord and peripheral blood and low expression in brain, skeletal muscle and spleen. Expressed also on Th2- and Tc2- type cells, eosinophils and basophils.


Volume
50 µl

GPR44 / CRTH2 Antibody (Extracellular Domain) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

GPR44 / CRTH2 Antibody (Extracellular Domain) - Images



Anti-GPR44 / CRTH2 antibody ALS10674 immunocytochemistry (ICC) staining of HEK293 human...

GPR44 / CRTH2 Antibody (Extracellular Domain) - Background

Receptor for prostaglandin D2 (PGD2). Coupled to the G(i)-protein. Receptor activation may result in pertussis toxin- sensitive decreases in cAMP levels and Ca(2+) mobilization. PI3K signaling is also implicated in mediating PTGDR2 effects. PGD2 induced receptor internalization. CRTH2 internalization can be regulated by diverse kinases such as, PKC, PKA, ADRBK1/GRK2, GPRK5/GRK5 and GRK6. Receptor activation is responsible, at least in part, in immune regulation and allergic/inflammation responses.

GPR44 / CRTH2 Antibody (Extracellular Domain) - References

Marchese A.,et al.Genomics 56:12-21(1999).
Nagata K.,et al.J. Immunol. 162:1278-1286(1999).
Methner A.,et al.Submitted (MAR-1999) to the EMBL/GenBank/DDBJ databases.
King M.M.,et al.Submitted (DEC-2003) to the EMBL/GenBank/DDBJ databases.
Taylor T.D.,et al.Nature 440:497-500(2006).