

GPRC5A / RAI3 Antibody (Transmembrane Domain)
Rabbit Polyclonal Antibody
Catalog # ALS10677**Specification**

GPRC5A / RAI3 Antibody (Transmembrane Domain) - Product Information

Application	IHC
Primary Accession	Q8NFJ5
Reactivity	Human, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40kDa KDa

GPRC5A / RAI3 Antibody (Transmembrane Domain) - Additional Information**Gene ID** 9052**Other Names**

Retinoic acid-induced protein 3, G-protein coupled receptor family C group 5 member A, Orphan G-protein-coupling receptor PEIG-1, Retinoic acid-induced gene 1 protein, RAIG-1, GPRC5A, GPCR5A, RAI3, RAIG1

Target/Specificity

Human GPRC5A / RAI3. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage

Long term: -70°C; Short term: +4°C

Precautions

GPRC5A / RAI3 Antibody (Transmembrane Domain) is for research use only and not for use in diagnostic or therapeutic procedures.

GPRC5A / RAI3 Antibody (Transmembrane Domain) - Protein Information**Name** GPRC5A**Synonyms** GPCR5A, RAI3, RAIG1**Function**

Orphan receptor. Could be involved in modulating differentiation and maintaining homeostasis of epithelial cells. This retinoic acid-inducible GPCR provide evidence for a possible interaction between retinoid and G-protein signaling pathways. Functions as a negative modulator of EGFR signaling (By similarity). May act as a lung tumor suppressor (PubMed:18000218).

Cellular Location

Cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle membrane; Multi-pass

membrane protein. Note=Localized in perinuclear vesicles, probably Golgi- associated vesicles.

Tissue Location

Expressed at high level in fetal and adult lung tissues but repressed in most human lung cancers (PubMed:9857033, PubMed:18000218). Constitutively expressed in fetal kidney and adult placenta, kidney, prostate, testis, ovary, small intestine, colon, stomach, and spinal cord at low to moderate levels. Not detectable in fetal heart, brain, and liver and adult heart, brain, liver, skeletal muscle, pancreas, spleen, thymus, and peripheral leukocytes. According to PubMed:10783259, expressed at low but detectable level in pancreas and heart.


Volume

50 μ l


GPRC5A / RAI3 Antibody (Transmembrane Domain) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GPRC5A / RAI3 Antibody (Transmembrane Domain) - Images


Anti-GPRC5A / RAI3 antibody IHC of human Pancreas, Carcinoma.




Anti-GPRC5A / RAI3 antibody IHC of human Skin, Melanoma.



Anti-GPRC5A / RAI3 antibody IHC of human Breast, Carcinoma.



Anti-GPRC5A / RAI3 antibody ALS10677 IHC of human breast.



Anti-GPRC5A / RAI3 antibody IHC of human testis.

GPRC5A / RAI3 Antibody (Transmembrane Domain) - Background

Unknown. This G-protein coupled receptor could be involved in modulating differentiation and maintaining homeostasis of epithelial cells. The comparable expression level in fetal lung and kidney with adult tissues suggests a possible role in embryonic development and maturation of these organs. This retinoic acid-inducible GPCR provide evidence for a possible interaction between retinoid and G-protein signaling pathways.

GPRC5A / RAI3 Antibody (Transmembrane Domain) - References

Cheng Y.,et al.J. Biol. Chem. 273:35008-35015(1998).
Cafferata E.G.,et al.Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Braeuner-Osborne H.,et al.Genomics 65:121-128(2000).