

DHRS7 Antibody (C-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS10953**Specification**

DHRS7 Antibody (C-Terminus) - Product Information

Application	IHC
Primary Accession	O9Y394
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38kDa KDa

DHRS7 Antibody (C-Terminus) - Additional Information**Gene ID** 51635**Other Names**

Dehydrogenase/reductase SDR family member 7, 1.1.-., Retinal short-chain dehydrogenase/reductase 4, retSDR4, DHRS7, DHRS7A, RETSDR4

Target/Specificity

Human DHRS7. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage

Long term: -70°C; Short term: +4°C

Precautions

DHRS7 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

DHRS7 Antibody (C-Terminus) - Protein Information**Name** DHRS7 ([HGNC:21524](#))**Function**

NADPH-dependent oxidoreductase which catalyzes the reduction of a variety of compounds bearing carbonyl groups including steroids, retinoids and xenobiotics (PubMed:24246760, PubMed:26466768, PubMed:28687384, PubMed:28457967). Catalyzes the reduction/inactivation of 5alpha-dihydrotestosterone to 3alpha-androstanediol, with a possible role in the modulation of androgen receptor function (PubMed:28687384, PubMed:28457967). Involved in the reduction of all-trans-retinal to all-trans-retinol (PubMed:28457967).

[26466768](http://www.uniprot.org/citations/26466768)). Converts cortisone to 20beta- dihydrocortisone in vitro, although the physiological relevance of this activity is questionable (PubMed: [28457967](http://www.uniprot.org/citations/28457967)). Reduces exogenous compounds such as quinones (1,2-naphthoquinone, 9,10-phenanthrenequinone and benzoquinone) and other xenobiotics (alpha-diketones) in vitro, suggesting a role in the biotransformation of xenobiotics with carbonyl group (PubMed: [24246760](http://www.uniprot.org/citations/24246760)), PubMed: [26466768](http://www.uniprot.org/citations/26466768)). A dehydrogenase activity has not been detected so far (PubMed: [24246760](http://www.uniprot.org/citations/24246760)). May play a role as tumor suppressor (PubMed: [26311046](http://www.uniprot.org/citations/26311046)).

Cellular Location

Endoplasmic reticulum membrane. Note=Bound to the endoplasmic reticulum membrane, possibly through a N-terminus anchor. The main bulk of the polypeptide chain was first reported to be facing toward the lumen of the endoplasmic reticulum (PubMed:24246760) However, it was later shown to be facing the cytosol (PubMed:28457967)

Tissue Location

Found predominantly in the adrenal glands, liver, thyroid, prostate, small intestine, colon, stomach, kidney and brain (PubMed:26466768). Lower levels observed in skeletal muscle, the lung and the spleen (PubMed:26466768).

Volume


50 µl


DHRS7 Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DHRS7 Antibody (C-Terminus) - Images





Anti-DHRS7 antibody ALS10953 IHC of human prostate.



Anti-DHRS7 antibody ALS10953 IHC of human kidney.

DHRS7 Antibody (C-Terminus) - References

- Lai C.-H., et al. *Genome Res.* 10:703-713(2000).
Haeseleer F., et al. *Methods Enzymol.* 316:372-383(2000).
Clark H.F., et al. *Genome Res.* 13:2265-2270(2003).
Ota T., et al. *Nat. Genet.* 36:40-45(2004).
Heilig R., et al. *Nature* 421:601-607(2003).