

VRL1 / TRPV2 Antibody (Internal)
Rabbit Polyclonal Antibody
Catalog # ALS11008**Specification**

VRL1 / TRPV2 Antibody (Internal) - Product Information

Application	IHC
Primary Accession	Q9Y5S1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	86kDa KDa

VRL1 / TRPV2 Antibody (Internal) - Additional Information**Gene ID** 51393**Other Names**

Transient receptor potential cation channel subfamily V member 2, TrpV2, Osm-9-like TRP channel 2, OTRPC2, Vanilloid receptor-like protein 1, VRL-1, TRPV2, VRL

Target/Specificity

Human TRPV2. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage

Long term: -70°C; Short term: +4°C

Precautions

VRL1 / TRPV2 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

VRL1 / TRPV2 Antibody (Internal) - Protein Information**Name** TRPV2**Synonyms** VRL**Function**

Calcium-permeable, non-selective cation channel with an outward rectification. Seems to be regulated, at least in part, by IGF- I, PDGF and neuropeptide head activator. May transduce physical stimuli in mast cells. Activated by temperatures higher than 52 degrees Celsius; is not activated by vanilloids and acidic pH.

Cellular Location

Cell membrane; Multi-pass membrane protein. Cytoplasm. Melanosome. Note=Translocates from the cytoplasm to the plasma membrane upon ligand stimulation (By similarity). Identified by mass spectrometry in melanosome fractions from stage I to stage IV


Volume
50 µl

VRL1 / TRPV2 Antibody (Internal) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

VRL1 / TRPV2 Antibody (Internal) - Images



Anti-TRPV2 antibody ALS11008 IHC of human prostate.



Anti-TRPV2 antibody ALS11008 IHC of human skeletal muscle.

VRL1 / TRPV2 Antibody (Internal) - Background

Calcium-permeable, non-selective cation channel with an outward rectification. Seems to be regulated, at least in part, by IGF-I, PDGF and neuropeptide head activator. May transduce physical stimuli in mast cells. Activated by temperatures higher than 52 degrees Celsius; is not activated by

vanilloids and acidic pH.

VRL1 / TRPV2 Antibody (Internal) - References

Caterina M.J.,et al.Nature 398:436-441(1999).

Garcia R.L.,et al.Submitted (NOV-1998) to the EMBL/GenBank/DDBJ databases.

Kelsell R.E.,et al.Submitted (MAY-2002) to the EMBL/GenBank/DDBJ databases.

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Zody M.C.,et al.Nature 440:1045-1049(2006).