

ITGA6/Integrin Alpha 6/CD49f Antibody (Internal)
Rabbit Polyclonal Antibody
Catalog # ALS11016**Specification**

ITGA6/Integrin Alpha 6/CD49f Antibody (Internal) - Product Information

Application	IHC
Primary Accession	P23229
Reactivity	Human, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	127kDa KDa

ITGA6/Integrin Alpha 6/CD49f Antibody (Internal) - Additional Information**Gene ID** 3655**Other Names**

Integrin alpha-6, CD49 antigen-like family member F, VLA-6, CD49f, Integrin alpha-6 heavy chain, Integrin alpha-6 light chain, Processed integrin alpha-6, Alpha6p, ITGA6

Target/Specificity

Human ITGA6 / CD49f. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage

Long term: -70°C; Short term: +4°C

Precautions

ITGA6/Integrin Alpha 6/CD49f Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

ITGA6/Integrin Alpha 6/CD49f Antibody (Internal) - Protein Information**Name** ITGA6**Function**

Integrin alpha-6/beta-1 (ITGA6:ITGB1) is a receptor for laminin on platelets (By similarity). Integrin alpha-6/beta-1 (ITGA6:ITGB1) is present in oocytes and is involved in sperm-egg fusion (By similarity). Integrin alpha-6/beta-4 (ITGA6:ITGB4) is a receptor for laminin in epithelial cells and it plays a critical structural role in the hemidesmosome (By similarity). ITGA6:ITGB4 binds to NRG1 (via EGF domain) and this binding is essential for NRG1-ERBB signaling (PubMed:20682778). ITGA6:ITGB4 binds to IGF1 and this binding is essential for IGF1 signaling (PubMed:22351760). ITGA6:ITGB4 binds to IGF2 and this binding is essential for IGF2 signaling (PubMed:28873464).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell membrane; Lipid-anchor

Tissue Location

Integrin alpha-6/beta-4 is predominantly expressed by epithelia. Isoforms containing segment X1 are ubiquitously expressed. Isoforms containing segment X1X2 are expressed in heart, kidney, placenta, colon, duodenum, myoblasts and myotubes, and in a limited number of cell lines; they are always coexpressed with the ubiquitous isoform containing segment X1. In some tissues (e.g Salivary gland), isoforms containing cytoplasmic segment A and isoforms containing segment B are detected while in others, only isoforms containing one cytoplasmic segment are found (segment A in epidermis and segment B in kidney). Processed integrin alpha-6: Expressed at low levels in normal prostate tissue with elevated levels in prostate cancer tissue (at protein level) (PubMed:15023541)


Volume

50 µl

ITGA6/Integrin Alpha 6/CD49f Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ITGA6/Integrin Alpha 6/CD49f Antibody (Internal) - Images

Anti-ITGA6 / CD49f antibody ALS11016 IHC of human breast.

ITGA6/Integrin Alpha 6/CD49f Antibody (Internal) - Background

Integrin alpha-6/beta-1 is a receptor for laminin on platelets. Integrin alpha-6/beta-4 is a receptor for laminin in epithelial cells and it plays a critical structural role in the hemidesmosome.

ITGA6/Integrin Alpha 6/CD49f Antibody (Internal) - References

Tamura R.N.,et al.J. Cell Biol. 111:1593-1604(1990).
Quaranta V.,et al.Submitted (JUN-1991) to the EMBL/GenBank/DDBJ databases.
Pulkkinen L.,et al.Submitted (JUL-1999) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Hillier L.W.,et al.Nature 434:724-731(2005).