10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

GCNT3 Antibody (Internal)
Rabbit Polyclonal Antibody
Catalog # ALS11096

Specification

GCNT3 Antibody (Internal) - Product Information

Application	**IHC**
Primary Accession	O95395
Reactivity	**Human, Horse, Dog**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**51kDa KDa**

GCNT3 Antibody (Internal) - Additional Information

Gene ID 9245

Other Names
Beta-1, 3-galactosyl-O-glycosyl-glycoprotein beta-1, 6-N-acetylglucosaminyltransferase 3, 2.4.1.102, 2.4.1.150, C2GnT-mucin type, C2GnT-M, hC2GnT-M, Core 2/core 4 beta-1, 6-N-acetylglucosaminyltransferase, C2/4GnT, GCNT3

Target/Specificity
Human GCNT3. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage
Long term: -70°C; Short term: +4°C

Precautions
GCNT3 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

GCNT3 Antibody (Internal) - Protein Information

Name GCNT3

Function
Glycosyltransferase that can synthesize all known mucin beta 6 N-acetylglucosaminides. Mediates core 2 and core 4 O-glycan branching, 2 important steps in mucin-type biosynthesis. Has also I-branching enzyme activity by converting linear into branched poly-N- acetyllactosaminoglycans, leading to introduce the blood group I antigen during embryonic development.

Cellular Location
Golgi apparatus membrane; Single- pass type II membrane protein

Tissue Location
Primarily expressed in mucus-secreting tissues. Expressed in colon, kidney, small intestine,

trachea, and stomach, where mucin is produced.


Volume
50 µl

GCNT3 Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

GCNT3 Antibody (Internal) - Images



Anti-GCNT3 antibody ALS11096 IHC of human stomach.

GCNT3 Antibody (Internal) - Background

Glycosyltransferase that can synthesize all known mucin beta 6 N-acetylglucosaminides. Mediates core 2 and core 4 O-glycan branching, 2 important steps in mucin-type biosynthesis. Has also I-branching enzyme activity by converting linear into branched poly-N-acetyllactosaminoglycans, leading to introduce the blood group I antigen during embryonic development.

GCNT3 Antibody (Internal) - References

Yeh J.-C.,et al.J. Biol. Chem. 274:3215-3221(1999).
Schwientek T.,et al.J. Biol. Chem. 274:4504-4512(1999).
Tan S.,et al.Am. J. Respir. Cell Mol. Biol. 36:737-745(2007).
Ishibashi Y.,et al.Glycoconj. J. 22:53-62(2005).
Huang M.-C.,et al.Oncogene 25:3267-3276(2006).