

RFFL Antibody (aa1-363)

Rabbit Polyclonal Antibody Catalog # ALS11353

Specification

RFFL Antibody (aa1-363) - Product Information

Application IHC
Primary Accession Q8WZ73
Reactivity Human
Host Rabbit
Clonality Polyclonal
Calculated MW 41kDa KDa

RFFL Antibody (aa1-363) - Additional Information

Gene ID 117584

Other Names

E3 ubiquitin-protein ligase rififylin, 6.3.2.-, Caspase regulator CARP2, Caspases-8 and -10-associated RING finger protein 2, CARP-2, FYVE-RING finger protein Sakura, Fring, RING finger and FYVE-like domain-containing protein 1, RING finger protein 189, RING finger protein 34-like, RFFL (HGNC:24821), RNF189, RNF34L

Target/Specificity

Amino acids 1-363 of human RFFL protein.

Reconstitution & Storage

+4°C or -20°C, Avoid repeated freezing and thawing.

Precautions

RFFL Antibody (aa1-363) is for research use only and not for use in diagnostic or therapeutic procedures.

RFFL Antibody (aa1-363) - Protein Information

Name RFFL (HGNC:24821)

Synonyms RNF189, RNF34L

Function

E3 ubiquitin-protein ligase that regulates several biological processes through the ubiquitin-mediated proteasomal degradation of various target proteins. Mediates 'Lys-48'-linked polyubiquitination of PRR5L and its subsequent proteasomal degradation thereby indirectly regulating cell migration through the mTORC2 complex. Ubiquitinates the caspases CASP8 and CASP10, promoting their proteasomal degradation, to negatively regulate cell death downstream of death domain receptors in the extrinsic pathway of apoptosis. Negatively regulates the tumor necrosis factor-mediated signaling pathway through targeting of RIPK1 to ubiquitin-mediated



proteasomal degradation. Negatively regulates p53/TP53 through its direct ubiquitination and targeting to proteasomal degradation. Indirectly, may also negatively regulate p53/TP53 through ubiquitination and degradation of SFN. May also play a role in endocytic recycling.

Cellular Location

Cytoplasm, cytosol. Cell membrane; Peripheral membrane protein. Recycling endosome membrane; Peripheral membrane protein. Note=The FYVE-type zinc finger may mediate phosphatidylinositol phosphate-binding and control subcellular localization

Tissue Location


Ubiquitous. Detected in spleen, thymus, prostate, testis, ovary, small intestine, colon and peripheral blood leukocytes

RFFL Antibody (aa1-363) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

RFFL Antibody (aa1-363) - Images



Anti-RFFL antibody IHC of human brain, cortex.

RFFL Antibody (aa1-363) - Background

E3 ubiquitin-protein ligase that regulates several biological processes through the ubiquitin-mediated proteasomal degradation of various target proteins. Mediates 'Lys-48'-linked polyubiquitination of PRR5L and its subsequent proteasomal degradation thereby indirectly regulating cell migration through the mTORC2 complex. Ubiquitinates the caspases CASP8 and CASP10, promoting their proteasomal degradation, to negatively regulate cell death downstream of death domain receptors in the extrinsic pathway of apoptosis. Negatively regulates the tumor necrosis factor-mediated signaling pathway through targeting of RIPK1 to ubiquitin-mediated proteasomal degradation. Negatively regulates p53/TP53 through its direct ubiquitination and targeting to proteasomal degradation. Indirectly, may also negatively regulate p53/TP53 through



ubiquitination and degradation of SFN. May also play a role in endocytic recycling.

RFFL Antibody (aa1-363) - References

Hong W.,et al.Submitted (OCT-2001) to the EMBL/GenBank/DDBJ databases. Kanbe D.,et al.Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases. Ota T.,et al.Nat. Genet. 36:40-45(2004). Bechtel S.,et al.BMC Genomics 8:399-399(2007). Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.