

TRAF2 Antibody (N-Terminus) Rabbit Polyclonal Antibody Catalog # ALS11734

## **Specification**

## **TRAF2 Antibody (N-Terminus) - Product Information**

Application Primary Accession Reactivity Host Clonality Calculated MW IHC <u>Q12933</u> Human, Mouse, Rat Rabbit Polyclonal 56kDa KDa

### TRAF2 Antibody (N-Terminus) - Additional Information

Gene ID 7186

**Other Names** TNF receptor-associated factor 2, 6.3.2.-, E3 ubiquitin-protein ligase TRAF2, Tumor necrosis factor type 2 receptor-associated protein 3, TRAF2, TRAP3

**Target/Specificity** 16 amino acid peptide from near the amino terminus of human TRAF2

**Reconstitution & Storage** Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

**Precautions** TRAF2 Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

### TRAF2 Antibody (N-Terminus) - Protein Information

Name TRAF2

Synonyms TRAP3

#### Function

Regulates activation of NF-kappa-B and JNK and plays a central role in the regulation of cell survival and apoptosis (PubMed:<a href="http://www.uniprot.org/citations/22212761" target="\_blank">22212761</a>). Required for normal antibody isotype switching from IgM to IgG. Has E3 ubiquitin-protein ligase activity and promotes 'Lys- 63'-linked ubiquitination of target proteins, such as BIRC3, RIPK1 and TICAM1. Is an essential constituent of several E3 ubiquitin-protein ligase complexes, where it promotes the ubiquitination of target proteins by bringing them into contact with other E3 ubiquitin ligases. Regulates BIRC2 and BIRC3 protein levels by inhibiting their autoubiquitination and subsequent degradation; this does not depend on the TRAF2 RING-type zinc finger domain. Plays a role in mediating activation of NF-kappa-B by EIF2AK2/PKR. In complex with BIRC2 or BIRC3, promotes ubiquitination of IKBKE.



Cellular Location Cytoplasm

## TRAF2 Antibody (N-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

### TRAF2 Antibody (N-Terminus) - Images



Anti-TRAF2 antibody IHC of human heart.



Anti-TRAF2 antibody IHC of human small intestine.

# TRAF2 Antibody (N-Terminus) - Background

Regulates activation of NF-kappa-B and JNK and plays a central role in the regulation of cell survival and apoptosis. Required for normal antibody isotype switching from IgM to IgG. Has E3 ubiquitin-protein ligase activity and promotes 'Lys-63'- linked ubiquitination of target proteins, such



as BIRC3, RIPK1 and TICAM1. Is an essential constituent of several E3 ubiquitin- protein ligase complexes, where it promotes the ubiquitination of target proteins by bringing them into contact with other E3 ubiquitin ligases. Regulates BIRC2 and BIRC3 protein levels by inhibiting their autoubiquitination and subsequent degradation; this does not depend on the TRAF2 RING-type zinc finger domain. Plays a role in mediating activation of NF-kappa-B by EIF2AK2/PKR. In complex with BIRC2 or BIRC3, promotes ubiquitination of IKBKE.

## TRAF2 Antibody (N-Terminus) - References

Song H.Y.,et al.Biochem. J. 309:825-829(1995). Ota T.,et al.Nat. Genet. 36:40-45(2004). Bechtel S.,et al.BMC Genomics 8:399-399(2007). Humphray S.J.,et al.Nature 429:369-374(2004). Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.