

ATM Antibody (aa2577-3056, clone 2C1)
Mouse Monoclonal Antibody
Catalog # ALS11851**Specification****ATM Antibody (aa2577-3056, clone 2C1) - Product Information**

Application	IHC
Primary Accession	Q13315
Reactivity	Human, Mouse, Rat, Monkey
Host	Mouse
Clonality	Monoclonal
Calculated MW	351kDa KDa

ATM Antibody (aa2577-3056, clone 2C1) - Additional Information**Gene ID** 472**Other Names**

Serine-protein kinase ATM, 2.7.11.1, Ataxia telangiectasia mutated, A-T mutated, ATM

Target/Specificity

Recombinant protein expressed in E. coli corresponding to amino acids 2577-3056.

Reconstitution & Storage

+4°C or -20°C, Avoid repeated freezing and thawing.

Precautions

ATM Antibody (aa2577-3056, clone 2C1) is for research use only and not for use in diagnostic or therapeutic procedures.

ATM Antibody (aa2577-3056, clone 2C1) - Protein Information**Name** ATM**Function**

Serine/threonine protein kinase which activates checkpoint signaling upon double strand breaks (DSBs), apoptosis and genotoxic stresses such as ionizing ultraviolet A light (UVA), thereby acting as a DNA damage sensor (PubMed:<[a href="http://www.uniprot.org/citations/9733514" target="_blank">9733514](http://www.uniprot.org/citations/9733514), PubMed:<[a href="http://www.uniprot.org/citations/10550055" target="_blank">10550055](http://www.uniprot.org/citations/10550055), PubMed:<[a href="http://www.uniprot.org/citations/10839545" target="_blank">10839545](http://www.uniprot.org/citations/10839545), PubMed:<[a href="http://www.uniprot.org/citations/10910365" target="_blank">10910365](http://www.uniprot.org/citations/10910365), PubMed:<[a href="http://www.uniprot.org/citations/12556884" target="_blank">12556884](http://www.uniprot.org/citations/12556884), PubMed:<[a href="http://www.uniprot.org/citations/14871926" target="_blank">14871926](http://www.uniprot.org/citations/14871926), PubMed:<[a href="http://www.uniprot.org/citations/15456891" target="_blank">15456891](http://www.uniprot.org/citations/15456891), PubMed:<[a href="http://www.uniprot.org/citations/15448695" target="_blank">15448695](http://www.uniprot.org/citations/15448695), PubMed:<[a href="http://www.uniprot.org/citations/15916964" target="_blank">15916964](http://www.uniprot.org/citations/15916964), PubMed:<[a href="http://www.uniprot.org/citations/17923702" target="_blank">17923702](http://www.uniprot.org/citations/17923702)). Recognizes the substrate consensus sequence [ST]-Q

(PubMed:9733514, PubMed:10550055, PubMed:10839545, PubMed:10910365, PubMed:12556884, PubMed:14871926, PubMed:15456891, PubMed:15448695, PubMed:15916964, PubMed:17923702). Phosphorylates 'Ser-139' of histone variant H2AX at double strand breaks (DSBs), thereby regulating DNA damage response mechanism (By similarity). Also plays a role in pre-B cell allelic exclusion, a process leading to expression of a single immunoglobulin heavy chain allele to enforce clonality and monospecific recognition by the B-cell antigen receptor (BCR) expressed on individual B-lymphocytes. After the introduction of DNA breaks by the RAG complex on one immunoglobulin allele, acts by mediating a repositioning of the second allele to pericentromeric heterochromatin, preventing accessibility to the RAG complex and recombination of the second allele. Also involved in signal transduction and cell cycle control. May function as a tumor suppressor. Necessary for activation of ABL1 and SAPK. Phosphorylates DYRK2, CHEK2, p53/TP53, FBXW7, FANCD2, NFKBIA, BRCA1, CTIP, nibrin (NBN), TERF1, UFL1, RAD9, UBQLN4 and DCLRE1C (PubMed:9843217, PubMed:9733515, PubMed:10550055, PubMed:10766245, PubMed:10839545, PubMed:10910365, PubMed:10802669, PubMed:10973490, PubMed:11375976, PubMed:12086603, PubMed:15456891, PubMed:19965871, PubMed:30612738, PubMed:30886146, PubMed:26774286). May play a role in vesicle and/or protein transport. Could play a role in T-cell development, gonad and neurological function. Plays a role in replication-dependent histone mRNA degradation. Binds DNA ends. Phosphorylation of DYRK2 in nucleus in response to genotoxic stress prevents its MDM2-mediated ubiquitination and subsequent proteasome degradation (PubMed:19965871). Phosphorylates ATF2 which stimulates its function in DNA damage response (PubMed:15916964). Phosphorylates ERCC6 which is essential for its chromatin remodeling activity at DNA double-strand breaks (PubMed:29203878). Phosphorylates TTC5/STRAP at 'Ser-203' in the cytoplasm in response to DNA damage, which promotes TTC5/STRAP nuclear localization (PubMed:15448695). Also involved in pexophagy by mediating phosphorylation of PEX5: translocated to peroxisomes in response to reactive oxygen species (ROS), and catalyzes phosphorylation of PEX5, promoting PEX5 ubiquitination and induction of pexophagy (PubMed:26344566).

Cellular Location

Nucleus. Cytoplasmic vesicle. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome {ECO:0000250|UniProtKB:Q62388}. Peroxisome matrix. Note=Primarily nuclear (PubMed:9050866, PubMed:9150358). Found also in endocytic vesicles in association with beta-adaptin (PubMed:9707615). Translocated to peroxisomes in response to reactive oxygen

species (ROS) by PEX5 (PubMed:26344566)

Tissue Location


Found in pancreas, kidney, skeletal muscle, liver, lung, placenta, brain, heart, spleen, thymus, testis, ovary, small intestine, colon and leukocytes

ATM Antibody (aa2577-3056, clone 2C1) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ATM Antibody (aa2577-3056, clone 2C1) - Images



Anti-ATM antibody IHC of human testis.

ATM Antibody (aa2577-3056, clone 2C1) - Background

Serine/threonine protein kinase which activates checkpoint signaling upon double strand breaks (DSBs), apoptosis and genotoxic stresses such as ionizing ultraviolet A light (UVA), thereby acting as a DNA damage sensor. Recognizes the substrate consensus sequence [ST]-Q. Phosphorylates 'Ser-139' of histone variant H2AX/H2AFX at double strand breaks (DSBs), thereby regulating DNA damage response mechanism. Also plays a role in pre-B cell allelic exclusion, a process leading to expression of a single immunoglobulin heavy chain allele to enforce clonality and monospecific recognition by the B-cell antigen receptor (BCR) expressed on individual B-lymphocytes. After the introduction of DNA breaks by the RAG complex on one immunoglobulin allele, acts by mediating a repositioning of the second allele to pericentromeric heterochromatin, preventing accessibility to the RAG complex and recombination of the second allele. Also involved in signal transduction and cell cycle control. May function as a tumor suppressor. Necessary for activation of ABL1 and SAPK. Phosphorylates DYRK2, CHEK2, p53/TP53, FANCD2, NFKBIA, BRCA1, CTIP, nibrin (NBN), TERF1, RAD9 and DCLRE1C. May play a role in vesicle and/or protein transport. Could play a role in T-cell development, gonad and neurological function. Plays a role in replication-dependent histone mRNA degradation. Binds DNA ends. Phosphorylation of DYRK2 in nucleus in response to genotoxic stress

prevents its MDM2-mediated ubiquitination and subsequent proteasome degradation.
Phosphorylates ATF2 which stimulates its function in DNA damage response.

ATM Antibody (aa2577-3056, clone 2C1) - References

Savitsky K.,et al.Hum. Mol. Genet. 4:2025-2032(1995).

Vorechovsky I.,et al.Cancer Res. 56:2726-2732(1996).

Platzer M.,et al.Genome Res. 7:592-605(1997).

Taylor T.D.,et al.Nature 440:497-500(2006).

Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.