10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CD59 Antibody (clone MEM-43)

Mouse Monoclonal Antibody
Catalog # ALS11973

Specification

CD59 Antibody (clone MEM-43) - Product Information

Application	**IHC**
Primary Accession	P13987
Reactivity	**Human**
Host	**Mouse**
Clonality	**Monoclonal**
Calculated MW	**14kDa KDa**

CD59 Antibody (clone MEM-43) - Additional Information

Gene ID 966

Other Names
CD59 glycoprotein, 1F5 antigen, 20 kDa homologous restriction factor, HRF-20, HRF20, MAC-inhibitory protein, MAC-IP, MEM43 antigen, Membrane attack complex inhibition factor, MACIF, Membrane inhibitor of reactive lysis, MIRL, Protectin, CD59, CD59, MIC11, MIN1, MIN2, MIN3, MSK21

Target/Specificity
Thymocytes and T lymphocytes

Reconstitution & Storage
Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.

Precautions
CD59 Antibody (clone MEM-43) is for research use only and not for use in diagnostic or therapeutic procedures.

CD59 Antibody (clone MEM-43) - Protein Information

Name CD59

Synonyms MIC11, MIN1, MIN2, MIN3, MSK21

Function
Potent inhibitor of the complement membrane attack complex (MAC) action. Acts by binding to the C8 and/or C9 complements of the assembling MAC, thereby preventing incorporation of the multiple copies of C9 required for complete formation of the osmolytic pore. This inhibitor appears to be species-specific. Involved in signal transduction for T-cell activation complexed to a protein tyrosine kinase.

Cellular Location
Cell membrane; Lipid-anchor, GPI-anchor. Secreted. Note=Soluble form found in a number of


tissues

CD59 Antibody (clone MEM-43) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CD59 Antibody (clone MEM-43) - Images



Anti-CD59 antibody IHC of human placenta.

CD59 Antibody (clone MEM-43) - Background

Potent inhibitor of the complement membrane attack complex (MAC) action. Acts by binding to the C8 and/or C9 complements of the assembling MAC, thereby preventing incorporation of the multiple copies of C9 required for complete formation of the osmolytic pore. This inhibitor appears to be species-specific. Involved in signal transduction for T-cell activation complexed to a protein tyrosine kinase.

CD59 Antibody (clone MEM-43) - References

Davies A.,et al.J. Exp. Med. 170:637-654(1989).
Philbrick W.M.,et al.Eur. J. Immunol. 20:87-92(1990).
Okada H.,et al.Biochem. Biophys. Res. Commun. 162:1553-1559(1989).
Sugita Y.,et al.J. Biochem. 106:555-557(1989).
Sawada R.,et al.DNA Cell Biol. 9:213-220(1990).