

DCAF6 / NRIP Antibody (aa800-850)
Rabbit Polyclonal Antibody
Catalog # ALS12089**Specification**

DCAF6 / NRIP Antibody (aa800-850) - Product Information

Application	WB, IHC
Primary Accession	Q58WW2
Reactivity	Human, Mouse, Rat, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	96kDa KDa

DCAF6 / NRIP Antibody (aa800-850) - Additional Information**Gene ID** 55827**Other Names**

DDB1- and CUL4-associated factor 6, Androgen receptor complex-associated protein, ARCAP, IQ motif and WD repeat-containing protein 1, Nuclear receptor interaction protein, NRIP, DCAF6, IQWD1

Target/Specificity

A portion of amino acids 800-850 of human NRIP was used as the immunogen.

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.

Precautions

DCAF6 / NRIP Antibody (aa800-850) is for research use only and not for use in diagnostic or therapeutic procedures.

DCAF6 / NRIP Antibody (aa800-850) - Protein Information**Name** DCAF6**Synonyms** IQWD1**Function**

Ligand-dependent coactivator of nuclear receptors. Enhance transcriptional activity of the nuclear receptors NR3C1 and AR. May function as a substrate receptor for CUL4-DDB1 E3 ubiquitin-protein ligase complex.

Cellular Location

Nucleus.

Tissue Location

Highly expressed in skeletal muscle and testis. Expressed to a lesser degree in heart, prostate,

and adrenal gland

DCAF6 / NRIP Antibody (aa800-850) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)


- [Flow Cytometry](#)

- [Cell Culture](#)

DCAF6 / NRIP Antibody (aa800-850) - Images



Western blot of NRIP in human muscle tissue lysate in the 1) absence and 2) presence of...



Anti-DCAF6 / NRIP antibody IHC of human skeletal muscle.

DCAF6 / NRIP Antibody (aa800-850) - Background

Ligand-dependent coactivator of nuclear receptors. Enhance transcriptional activity of the nuclear receptors NR3C1 and AR. May function as a substrate receptor for CUL4-DDB1 E3 ubiquitin-protein ligase complex.

DCAF6 / NRIP Antibody (aa800-850) - References

Tsai T.C.,et al.J. Biol. Chem. 280:20000-20009(2005).
Chang T.-J.,et al.Submitted (JUN-2006) to the EMBL/GenBank/DDBJ databases.
Zhao B.,et al.Submitted (DEC-1998) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Gregory S.G.,et al.Nature 441:315-321(2006).