10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

SET / TAF-I Antibody (aa66-81)

Rabbit Polyclonal Antibody
Catalog # ALS12100

Specification

SET / TAF-I Antibody (aa66-81) - Product Information

Application	**WB, IHC**
Primary Accession	Q01105
Reactivity	**Human, Mouse, Rat, Zebrafish, Hamster, Monkey, Pig, Chicken, Horse, Xenopus, Bovine**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**33kDa KDa**

SET / TAF-I Antibody (aa66-81) - Additional Information

Gene ID 6418

Other Names
Protein SET, HLA-DR-associated protein II, Inhibitor of granzyme A-activated DNase, IGAAD, PHAPII, Phosphatase 2A inhibitor I2PP2A, I-2PP2A, Template-activating factor I, TAF-I, SET

Target/Specificity
Synthetic peptides of amino acids 66-81 and 135-151 of human SET (Q01105). Sequence homology is 100% in mouse and rat.

Reconstitution & Storage
Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.

Precautions
SET / TAF-I Antibody (aa66-81) is for research use only and not for use in diagnostic or therapeutic procedures.

SET / TAF-I Antibody (aa66-81) - Protein Information

Name SET

Function
Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone chaperoning. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. In the course of cytotoxic T-lymphocyte (CTL)-induced apoptosis, GZMA cleaves SET, disrupting its binding to NME1 and releasing NME1 inhibition. Isoform 1 and isoform 2 are potent inhibitors of protein phosphatase 2A. Isoform 1 and isoform 2 inhibit EP300/CREBBP and PCAF- mediated acetylation of histones (HAT) and nucleosomes, most probably by masking the accessibility of lysines of histones to the acetylases. The predominant target for inhibition is histone H4. HAT inhibition leads to silencing of HAT-dependent transcription and prevents active demethylation of DNA. Both isoforms stimulate DNA replication of the adenovirus genome

complexed with viral core proteins; however, isoform 2 specific activity is higher.

Cellular Location
Cytoplasm, cytosol. Endoplasmic reticulum. Nucleus, nucleoplasm. Note=In the cytoplasm, found both in the cytosol and associated with the endoplasmic reticulum. The SET complex is associated with the endoplasmic reticulum. Following CTL attack and cleavage by GZMA, moves rapidly to the nucleus, where it is found in the nucleoplasm, avoiding the nucleolus. Similar translocation to the nucleus is also observed for lymphocyte-activated killer cells after the addition of calcium


Tissue Location
Widely expressed. Low levels in quiescent cells during serum starvation, contact inhibition or differentiation. Highly expressed in Wilms' tumor

SET / TAF-I Antibody (aa66-81) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

SET / TAF-I Antibody (aa66-81) - Images



Western blot of INHAT-1 using antibody against Jurkat cell lysate.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Anti-SET antibody IHC of human skin.

SET / TAF-I Antibody (aa66-81) - Background

Multitasking protein, involved in apoptosis, transcription, nucleosome assembly and histone chaperoning. Isoform 2 anti-apoptotic activity is mediated by inhibition of the GZMA-activated DNase, NME1. In the course of cytotoxic T- lymphocyte (CTL)-induced apoptosis, GZMA cleaves SET, disrupting its binding to NME1 and releasing NME1 inhibition. Isoform 1 and isoform 2 are potent inhibitors of protein phosphatase 2A. Isoform 1 and isoform 2 inhibit EP300/CREBBP and PCAF-mediated acetylation of histones (HAT) and nucleosomes, most probably by masking the accessibility of lysines of histones to the acetylases. The predominant target for inhibition is histone H4. HAT inhibition leads to silencing of HAT-dependent transcription and prevents active demethylation of DNA. Both isoforms stimulate DNA replication of the adenovirus genome complexed with viral core proteins; however, isoform 2 specific activity is higher.

SET / TAF-I Antibody (aa66-81) - References

von Lindern M.,et al.Mol. Cell. Biol. 12:3346-3355(1992).
Vaesen M.,et al.Biol. Chem. Hoppe-Seyler 375:113-126(1994).
Nagata K.,et al.Proc. Natl. Acad. Sci. U.S.A. 92:4279-4283(1995).
Li M.,et al.J. Biol. Chem. 271:11059-11062(1996).
Tsujio I.,et al.FEBS Lett. 579:363-372(2005).