

I-FABP / FABP2 Antibody (C-Terminus)
Goat Polyclonal Antibody
Catalog # ALS12247**Specification**

I-FABP / FABP2 Antibody (C-Terminus) - Product Information

Application	IHC
Primary Accession	P12104
Reactivity	Human, Monkey
Host	Goat
Clonality	Polyclonal
Calculated MW	15kDa KDa

I-FABP / FABP2 Antibody (C-Terminus) - Additional Information**Gene ID** 2169**Other Names**

Fatty acid-binding protein, intestinal, Fatty acid-binding protein 2, Intestinal-type fatty acid-binding protein, I-FABP, FABP2, FABPI

Target/Specificity

Human FABP2.

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

I-FABP / FABP2 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

I-FABP / FABP2 Antibody (C-Terminus) - Protein Information**Name** FABP2**Synonyms** FABPI**Function**

FABPs are thought to play a role in the intracellular transport of long-chain fatty acids and their acyl-CoA esters. FABP2 is probably involved in triglyceride-rich lipoprotein synthesis. Binds saturated long-chain fatty acids with a high affinity, but binds with a lower affinity to unsaturated long-chain fatty acids. FABP2 may also help maintain energy homeostasis by functioning as a lipid sensor.

Cellular Location

Cytoplasm.

Tissue Location


Expressed in the small intestine and at much lower levels in the large intestine. Highest expression levels in the jejunum.

I-FABP / FABP2 Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

I-FABP / FABP2 Antibody (C-Terminus) - Images



Anti-FABP2 antibody IHC of human small intestine.

I-FABP / FABP2 Antibody (C-Terminus) - Background

FABP are thought to play a role in the intracellular transport of long-chain fatty acids and their acyl-CoA esters. FABP2 is probably involved in triglyceride-rich lipoprotein synthesis. Binds saturated long-chain fatty acids with a high affinity, but binds with a lower affinity to unsaturated long-chain fatty acids. FABP2 may also help maintain energy homeostasis by functioning as a lipid sensor.

I-FABP / FABP2 Antibody (C-Terminus) - References

Sweetser D.A., et al. J. Biol. Chem. 262:16060-16071(1987).
Hillier L.W., et al. Nature 434:724-731(2005).
Pelsers M.M.A.L., et al. Clin. Biochem. 36:529-535(2003).
Darimont C., et al. Am. J. Physiol. 276:G606-G612(1999).
Rajabzadeh M., et al. Biochemistry 42:12192-12199(2003).