10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ALDH1A1 / ALDH1 Antibody (Internal)

Goat Polyclonal Antibody
Catalog # ALS12289

Specification

ALDH1A1 / ALDH1 Antibody (Internal) - Product Information

Application	**IHC**
Primary Accession	P00352
Reactivity	**Human, Sheep, Bovine**
Host	**Goat**
Clonality	**Polyclonal**
Calculated MW	**55kDa KDa**

ALDH1A1 / ALDH1 Antibody (Internal) - Additional Information

Gene ID 216

Other Names
Retinal dehydrogenase 1, RALDH 1, RalDH1, 1.2.1.36, ALDH-E1, ALHDII, Aldehyde dehydrogenase family 1 member A1, Aldehyde dehydrogenase, cytosolic, ALDH1A1, ALDC, ALDH1, PUMB1

Target/Specificity
Human ALDH1A1.

Reconstitution & Storage
Store at -20°C. Minimize freezing and thawing.

Precautions
ALDH1A1 / ALDH1 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

ALDH1A1 / ALDH1 Antibody (Internal) - Protein Information

Name ALDH1A1 (HGNC:402)

Function
Cytosolic dehydrogenase that catalyzes the irreversible oxidation of a wide range of aldehydes to their corresponding carboxylic acid (PubMed:19296407, PubMed:12941160, PubMed:15623782, PubMed:17175089, PubMed:26373694, PubMed:25450233). Functions downstream of retinol dehydrogenases and catalyzes the oxidation of retinaldehyde into retinoic acid, the second step in the oxidation of retinol/vitamin A into retinoic acid (By similarity). This pathway is crucial to control the levels of retinol and retinoic acid, two important molecules which excess can be teratogenic and cytotoxic (By similarity). Also oxidizes aldehydes resulting from lipid peroxidation like

(E)-4-hydroxynon-2-enal/HNE, malonaldehyde and hexanal that form protein adducts and are highly cytotoxic. By participating for instance to the clearance of (E)-4-hydroxynon-2-enal/HNE in the lens epithelium prevents the formation of HNE-protein adducts and lens opacification (PubMed:19296407, PubMed:12941160, PubMed:15623782). Functions also downstream of fructosamine-3-kinase in the fructosamine degradation pathway by catalyzing the oxidation of 3-deoxyglucosone, the carbohydrate product of fructosamine 3-phosphate decomposition, which is itself a potent glycating agent that may react with lysine and arginine side-chains of proteins (PubMed:17175089). Has also an aminobutyraldehyde dehydrogenase activity and is probably part of an alternative pathway for the biosynthesis of GABA/4-aminobutanoate in midbrain, thereby playing a role in GABAergic synaptic transmission (By similarity).

Cellular Location
Cytoplasm, cytosol. Cell projection, axon {ECO:0000250|UniProtKB:P24549}


Tissue Location
Expressed by erythrocytes (at protein level).

ALDH1A1 / ALDH1 Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ALDH1A1 / ALDH1 Antibody (Internal) - Images



Anti-ALDH1A1 antibody IHC of human testis.

ALDH1A1 / ALDH1 Antibody (Internal) - Background

Binds free retinal and cellular retinol-binding protein- bound retinal. Can convert/oxidize retinaldehyde to retinoic acid (By similarity).

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ALDH1A1 / ALDH1 Antibody (Internal) - References

Hsu L.C.,et al.Genomics 5:857-865(1989).
Zheng C.F.,et al.Alcohol. Clin. Exp. Res. 17:828-831(1993).
Ramana K.V.,et al.Submitted (SEP-2003) to the EMBL/GenBank/DDBJ databases.
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Humphray S.J.,et al.Nature 429:369-374(2004).