

NT5E / eNT / CD73 Antibody (aa27-252, clone 2B6)
Mouse Monoclonal Antibody
Catalog # ALS12559**Specification****NT5E / eNT / CD73 Antibody (aa27-252, clone 2B6) - Product Information**

Application	IHC
Primary Accession	P21589
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	63kDa KDa

NT5E / eNT / CD73 Antibody (aa27-252, clone 2B6) - Additional Information**Gene ID** 4907**Other Names**

5'-nucleotidase, 5'-NT, 3.1.3.5, Ecto-5'-nucleotidase, CD73, NT5E, NT5, NTE

Reconstitution & Storage

Long term: -20°C; Short term: +4°C; Avoid freeze-thaw cycles.

Precautions

NT5E / eNT / CD73 Antibody (aa27-252, clone 2B6) is for research use only and not for use in diagnostic or therapeutic procedures.

NT5E / eNT / CD73 Antibody (aa27-252, clone 2B6) - Protein Information**Name** NT5E**Synonyms** NT5, NTE**Function**

Catalyzes the hydrolysis of nucleotide monophosphates, releasing inorganic phosphate and the corresponding nucleoside, with AMP being the preferred substrate (PubMed:21933152, PubMed:34403084, PubMed:24887587, PubMed:22997138, PubMed:23142347). Shows a preference for ribonucleotide monophosphates over their equivalent deoxyribose forms (PubMed:34403084). Other substrates include IMP, UMP, GMP, CMP, dAMP, dCMP, dTMP, NAD and NMN (PubMed:21933152, PubMed:34403084, PubMed:24887587, PubMed:22997138, PubMed:23142347).

href="http://www.uniprot.org/citations/23142347" target="_blank">23142347).


Cellular Location

Cell membrane; Lipid-anchor, GPI-anchor


NT5E / eNT / CD73 Antibody (aa27-252, clone 2B6) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NT5E / eNT / CD73 Antibody (aa27-252, clone 2B6) - Images

Anti-NT5E / CD73 antibody IHC of human adrenal.



Anti-NT5E / CD73 antibody IHC of human pancreas.

NT5E / eNT / CD73 Antibody (aa27-252, clone 2B6) - Background

Hydrolyzes extracellular nucleotides into membrane permeable nucleosides. Exhibits AMP-, NAD-,

and NMN-nucleosidase activities.

NT5E / eNT / CD73 Antibody (aa27-252, clone 2B6) - References

Misumi Y.,et al.Eur. J. Biochem. 191:563-569(1990).

Otsuki T.,et al.DNA Res. 12:117-126(2005).

Mungall A.J.,et al.Nature 425:805-811(2003).

Hansen K.R.,et al.Gene 167:307-312(1995).

Zanoni L.,et al.Submitted (MAY-1998) to the EMBL/GenBank/DDBJ databases.