

### PDCD4 Antibody (aa1-469, clone K4C1)

Mouse Monoclonal Antibody
Catalog # ALS12564

#### **Specification**

### PDCD4 Antibody (aa1-469, clone K4C1) - Product Information

Application IHC, WB
Primary Accession Q53EL6
Reactivity Human
Host Mouse
Clonality Monoclonal
Calculated MW 52kDa KDa

#### PDCD4 Antibody (aa1-469, clone K4C1) - Additional Information

Gene ID 27250

#### **Other Names**

Programmed cell death protein 4, Neoplastic transformation inhibitor protein, Nuclear antigen H731-like, Protein 197/15a, PDCD4, H731

#### **Reconstitution & Storage**

Long term: -20°C; Short term: +4°C; Avoid freeze-thaw cycles.

#### **Precautions**

PDCD4 Antibody (aa1-469, clone K4C1) is for research use only and not for use in diagnostic or therapeutic procedures.

# PDCD4 Antibody (aa1-469, clone K4C1) - Protein Information

#### Name PDCD4

Synonyms H731

#### **Function**

Inhibits translation initiation and cap-dependent translation. May excert its function by hindering the interaction between EIF4A1 and EIF4G. Inhibits the helicase activity of EIF4A. Modulates the activation of JUN kinase. Down-regulates the expression of MAP4K1, thus inhibiting events important in driving invasion, namely, MAPK85 activation and consequent JUN-dependent transcription. May play a role in apoptosis. Tumor suppressor. Inhibits tumor promoter-induced neoplastic transformation. Binds RNA (By similarity).

#### **Cellular Location**

Nucleus {ECO:0000250|UniProtKB:Q61823}. Cytoplasm {ECO:0000250|UniProtKB:Q61823}. Note=Shuttles between the nucleus and cytoplasm (By similarity). Predominantly nuclear under normal growth conditions, and when phosphorylated at Ser-457 (PubMed:16357133)

### **Tissue Location**




Up-regulated in proliferative cells. Highly expressed in epithelial cells of the mammary gland. Reduced expression in lung cancer and colon carcinoma.

## PDCD4 Antibody (aa1-469, clone K4C1) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# PDCD4 Antibody (aa1-469, clone K4C1) - Images




Anti-PDCD4 antibody IHC of human breast.



Anti-PDCD4 antibody IHC of human spleen.





Cell lysates of HepG2(40 ug) were resolved by SDS-PAGE, transferred to NC membrane and probed...

### PDCD4 Antibody (aa1-469, clone K4C1) - Background

Inhibits translation initiation and cap-dependent translation. May excert its function by hindering the interaction between EIF4A1 and EIF4G. Inhibits the helicase activity of EIF4A. Modulates the activation of JUN kinase. Down-regulates the expression of MAP4K1, thus inhibiting events important in driving invasion, namely, MAPK85 activation and consequent JUN-dependent transcription. May play a role in apoptosis. Tumor suppressor. Inhibits tumor promoter-induced neoplastic transformation. Binds RNA (By similarity).

# PDCD4 Antibody (aa1-469, clone K4C1) - References

Matsuhashi S.,et al.Res. Commun. Biochem. Cell Mol. Biol. 1:109-120(1997). Azzoni L.,et al.J. Immunol. 161:3493-3500(1998). Ota T.,et al.Nat. Genet. 36:40-45(2004). Totoki Y.,et al.Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases. Deloukas P.,et al.Nature 429:375-381(2004).