10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

VAV3 Antibody (Internal)
Goat Polyclonal Antibody
Catalog # ALS12714

Specification

VAV3 Antibody (Internal) - Product Information

Application	**IHC**
Primary Accession	Q9UKW4
Reactivity	**Human, Monkey**
Host	**Goat**
Clonality	**Polyclonal**
Calculated MW	**98kDa KDa**

VAV3 Antibody (Internal) - Additional Information

Gene ID 10451

Other Names
Guanine nucleotide exchange factor VAV3, VAV-3, VAV3

Target/Specificity
Human VAV3. This antibody is expected to recognize both reported isoforms (NP_006104.4; NP_001073343.1).

Reconstitution & Storage
Store at -20°C. Minimize freezing and thawing.

Precautions
VAV3 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

VAV3 Antibody (Internal) - Protein Information

Name VAV3

Function
Exchange factor for GTP-binding proteins RhoA, RhoG and, to a lesser extent, Rac1. Binds physically to the nucleotide-free states of those GTPases. Plays an important role in angiogenesis. Its recruitment by phosphorylated EPHA2 is critical for EFNA1-induced RAC1 GTPase activation and vascular endothelial cell migration and assembly (By similarity). May be important for integrin-mediated signaling, at least in some cell types. In osteoclasts, along with SYK tyrosine kinase, required for signaling through integrin alpha-v/beta-1 (ITAGV-ITGB1), a crucial event for osteoclast proper cytoskeleton organization and function. This signaling pathway involves RAC1, but not RHO, activation. Necessary for proper wound healing. In the course of wound healing, required for the phagocytotic cup formation preceding macrophage phagocytosis of apoptotic neutrophils. Responsible for integrin beta-2 (ITGB2)-mediated macrophage adhesion and, to a lesser extent, contributes to beta-3 (ITGB3)-mediated adhesion. Does not affect integrin beta-1 (ITGB1)-mediated adhesion (By similarity).


Tissue Location
Isoform 1 and isoform 3 are widely expressed; both are expressed at very low levels in skeletal muscle. In keratinocytes, isoform 1 is less abundant than isoform 3. Isoform 3 is detected at very low levels, if any, in adrenal gland, bone marrow, spleen, fetal brain and spinal chord; in these tissues, isoform 1 is readily detectable.

VAV3 Antibody (Internal) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

VAV3 Antibody (Internal) - Images



Anti-VAV3 antibody IHC of human skin.



Anti-VAV3 antibody IHC of human spleen.

VAV3 Antibody (Internal) - Background

Exchange factor for GTP-binding proteins RhoA, RhoG and, to a lesser extent, Rac1. Binds physically to the nucleotide-free states of those GTPases. Plays an important role in angiogenesis. Its recruitment by phosphorylated EPHA2 is critical for EFNA1- induced RAC1 GTPase activation and vascular endothelial cell migration and assembly (By similarity). May be important for integrin-mediated signaling, at least in some cell types. In osteoclasts, along with SYK tyrosine kinase, required for signaling through integrin alpha-v/beta-1 (ITAGV-ITGB1), a crucial event for osteoclast proper cytoskeleton organization and function. This signaling pathway involves RAC1, but not RHO, activation. Necessary for proper wound healing. In the course of wound healing, required for the phagocytotic cup formation preceding macrophage phagocytosis of apoptotic neutrophils. Responsible for integrin beta-2 (ITGB2)-mediated macrophage adhesion and, to a lesser extent, contributes to beta-3 (ITGB3)- mediated adhesion. Does not affect integrin beta-1 (ITGB1)- mediated adhesion (By similarity).

VAV3 Antibody (Internal) - References

Trenkle T.,et al.Nucleic Acids Res. 26:3883-3891(1998).
Movilla N.,et al.Mol. Cell. Biol. 19:7870-7885(1999).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Gregory S.G.,et al.Nature 441:315-321(2006).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.