10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

EZR / Ezrin Antibody (Internal)
Goat Polyclonal Antibody
Catalog # ALS12719

Specification

EZR / Ezrin Antibody (Internal) - Product Information

Application	**IHC**
Primary Accession	P15311
Reactivity	**Human, Monkey**
Host	**Goat**
Clonality	**Polyclonal**
Calculated MW	**69kDa KDa**

EZR / Ezrin Antibody (Internal) - Additional Information

Gene ID 7430

Other Names
Ezrin, Cytovillin, Villin-2, p81, EZR, VIL2

Target/Specificity
Human EZR / Ezrin.

Reconstitution & Storage
Store at -20°C. Minimize freezing and thawing.

Precautions
EZR / Ezrin Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

EZR / Ezrin Antibody (Internal) - Protein Information

Name EZR

Synonyms VIL2

Function
Probably involved in connections of major cytoskeletal structures to the plasma membrane. In epithelial cells, required for the formation of microvilli and membrane ruffles on the apical pole. Along with PLEKHG6, required for normal macropinocytosis.

Cellular Location
Apical cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection. Cell projection, microvillus membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, ruffle membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cell projection, microvillus {ECO:0000250|UniProtKB:P26040}. Note=Localization to the apical membrane of parietal cells depends on the interaction with PALS1 Localizes to cell extensions and peripheral processes of astrocytes (By similarity). Microvillar

peripheral membrane protein (cytoplasmic side). {ECO:0000250|UniProtKB:P31977}


Tissue Location
Expressed in cerebral cortex, basal ganglia, hippocampus, hypophysis, and optic nerve. Weakly expressed in brain stem and diencephalon. Stronger expression was detected in gray matter of frontal lobe compared to white matter (at protein level). Component of the microvilli of intestinal epithelial cells. Preferentially expressed in astrocytes of hippocampus, frontal cortex, thalamus, parahippocampal cortex, amygdala, insula, and corpus callosum. Not detected in neurons in most tissues studied

EZR / Ezrin Antibody (Internal) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

EZR / Ezrin Antibody (Internal) - Images



Anti-EZR / Ezrin antibody IHC of human colon.



10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-EZR / Ezrin antibody IHC of human kidney.

EZR / Ezrin Antibody (Internal) - Background

Probably involved in connections of major cytoskeletal structures to the plasma membrane. In epithelial cells, required for the formation of microvilli and membrane ruffles on the apical pole. Along with PLEKHG6, required for normal macropinocytosis.

EZR / Ezrin Antibody (Internal) - References

Gould K.L.,et al.EMBO J. 8:4133-4142(1989).
Turunen O.,et al.J. Biol. Chem. 264:16727-16732(1989).
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Mungall A.J.,et al.Nature 425:805-811(2003).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.