10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

FASLG / Fas Ligand Antibody (N-Terminus)

Rabbit Polyclonal Antibody

Catalog # ALS12766

Specification

FASLG / Fas Ligand Antibody (N-Terminus) - Product Information

Application	**IHC**
Primary Accession	P48023
Reactivity	**Human, Mouse, Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**31kDa KDa**

FASLG / Fas Ligand Antibody (N-Terminus) - Additional Information

Gene ID 356

Other Names
Tumor necrosis factor ligand superfamily member 6, Apoptosis antigen ligand, APTL, CD95 ligand, CD95-L, Fas antigen ligand, Fas ligand, FasL, CD178, Tumor necrosis factor ligand superfamily member 6, membrane form, Tumor necrosis factor ligand superfamily member 6, soluble form, Receptor-binding FasL ectodomain, Soluble Fas ligand, sFasL, ADAM10-processed FasL form, APL, FasL intracellular domain, FasL ICD, SPPL2A-processed FasL form, SPA, FASLG, APT1LG1, CD95L, FASL, TNFSF6

Reconstitution & Storage
Store at 2°C to 8°C degrees. Do not freeze.

Precautions
FASLG / Fas Ligand Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

FASLG / Fas Ligand Antibody (N-Terminus) - Protein Information

Name FASLG

Synonyms APT1LG1, CD95L, FASL, TNFSF6

Function
Cytokine that binds to TNFRSF6/FAS, a receptor that transduces the apoptotic signal into cells (PubMed:26334989, PubMed:9228058). Involved in cytotoxic T-cell-mediated apoptosis, natural killer cell-mediated apoptosis and in T-cell development (PubMed:9228058, PubMed:7528780, PubMed:9427603). Initiates fratricidal/suicidal activation-induced cell death (AICD) in antigen- activated T-cells contributing to the termination of immune responses (By similarity).

TNFRSF6/FAS-mediated apoptosis has also a role in the induction of peripheral tolerance (By similarity). Binds to TNFRSF6B/DcR3, a decoy receptor that blocks apoptosis (PubMed:27806260).

Cellular Location
Cell membrane; Single-pass type II membrane protein. Cytoplasmic vesicle lumen Lysosome lumen. Note=Is internalized into multivesicular bodies of secretory lysosomes after phosphorylation by FGR and monoubiquitination (PubMed:17164290). Colocalizes with the SPPL2A protease at the cell membrane (PubMed:17557115) [FasL intracellular domain]: Nucleus. Note=The FasL ICD cytoplasmic form is translocated into the nucleus.


Volume
250 µl

FASLG / Fas Ligand Antibody (N-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

FASLG / Fas Ligand Antibody (N-Terminus) - Images



Anti-FASLG / FASL antibody IHC of human kidney.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Anti-FASLG / FASL antibody IHC of human spleen.

FASLG / Fas Ligand Antibody (N-Terminus) - Background

Cytokine that binds to TNFRSF6/FAS, a receptor that transduces the apoptotic signal into cells. May be involved in cytotoxic T-cell mediated apoptosis and in T-cell development. TNFRSF6/FAS-mediated apoptosis may have a role in the induction of peripheral tolerance, in the antigen-stimulated suicide of mature T-cells, or both. Binding to the decoy receptor TNFRSF6B/DcR3 modulates its effects.

FASLG / Fas Ligand Antibody (N-Terminus) - References

Alderson M.,et al.J. Exp. Med. 181:71-77(1995).
Takahashi T.,et al.Int. Immunol. 6:1567-1574(1994).
Schaetzlein C.E.,et al.Submitted (JUN-1995) to the EMBL/GenBank/DDBJ databases.
Mita E.,et al.Biochem. Biophys. Res. Commun. 204:468-474(1994).
Zeytun A.,et al.Submitted (JUL-2000) to the EMBL/GenBank/DDBJ databases.