

### **CUEDC2 Antibody (C-Terminus)**

Rabbit Polyclonal Antibody Catalog # ALS12994

# **Specification**

# **CUEDC2 Antibody (C-Terminus) - Product Information**

Application WB, ICC Primary Accession Q9H467

Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 32kDa KDa

# **CUEDC2 Antibody (C-Terminus) - Additional Information**

**Gene ID** 79004

#### **Other Names**

CUE domain-containing protein 2, CUEDC2, C10orf66

### **Reconstitution & Storage**

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

#### **Precautions**

CUEDC2 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

# **CUEDC2 Antibody (C-Terminus) - Protein Information**

#### Name CUEDC2

### Synonyms C10orf66

### **Function**

Down-regulates ESR1 protein levels through the ubiquitination-proteasome pathway, regardless of the presence of 17 beta-estradiol. Also involved in 17 beta-estradiol-induced ESR1 degradation. Controls PGR protein levels through a similar mechanism.

#### **Cellular Location**

Cytoplasm. Nucleus

#### **Tissue Location**

Significantly up-regulated in breast tumor tissues compared with matched adjacent normal tissues (at protein level) Levels inversely correlate with ESR1 in breast cancers and are lower in low-grade tumors compared to high-grade tumors




TGI. 000.070.1300 Fdx. 030.073.1300

# **CUEDC2 Antibody (C-Terminus) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# CUEDC2 Antibody (C-Terminus) - Images



Western blot of CUEDC2 in HeLa cell lysate with CUEDC2 antibody at (A) 1 and (B) 2 ug/ml.



Immunocytochemistry in HeLa cells with CUEDC2 antibody at 5 µg/ml.

# **CUEDC2 Antibody (C-Terminus) - Background**

Down-regulates ESR1 protein levels through the ubiquitination-proteasome pathway, regardless of the presence of 17 beta-estradiol. Also involved in 17 beta-estradiol-induced ESR1 degradation. Controls PGR protein levels through a similar mechanism.

#### **CUEDC2 Antibody (C-Terminus) - References**

Ikeda A., et al. Submitted (MAR-1999) to the EMBL/GenBank/DDBJ databases. Matsumoto K., et al. Submitted (AUG-2005) to the EMBL/GenBank/DDBJ databases. Deloukas P., et al. Nature 429:375-381(2004). Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases. Zhang P.-J., et al. EMBO J. 26:1831-1842(2007).