

**KRT17 / CK17 / Cytokeratin 17 Antibody (C-Terminus, clone V21-R)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS13063****Specification**

---

**KRT17 / CK17 / Cytokeratin 17 Antibody (C-Terminus, clone V21-R) - Product Information**

Application	IHC
Primary Accession	<a href="#">Q04695</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48kDa KDa

**KRT17 / CK17 / Cytokeratin 17 Antibody (C-Terminus, clone V21-R) - Additional Information****Gene ID** 3872**Other Names**

Keratin, type I cytoskeletal 17, 39.1, Cytokeratin-17, CK-17, Keratin-17, K17, KRT17

**Target/Specificity**

Human Cytokeratin 17

**Reconstitution & Storage**

Store at -20°C. Aliquot to avoid freeze/thaw cycles.

**Precautions**

KRT17 / CK17 / Cytokeratin 17 Antibody (C-Terminus, clone V21-R) is for research use only and not for use in diagnostic or therapeutic procedures.

**KRT17 / CK17 / Cytokeratin 17 Antibody (C-Terminus, clone V21-R) - Protein Information****Name** KRT17**Function**

Type I keratin involved in the formation and maintenance of various skin appendages, specifically in determining shape and orientation of hair (By similarity). Required for the correct growth of hair follicles, in particular for the persistence of the anagen (growth) state (By similarity). Modulates the function of TNF-alpha in the specific context of hair cycling. Regulates protein synthesis and epithelial cell growth through binding to the adapter protein SFN and by stimulating Akt/mTOR pathway (By similarity). Involved in tissue repair. May be a marker of basal cell differentiation in complex epithelia and therefore indicative of a certain type of epithelial 'stem cells'. Acts as a promoter of epithelial proliferation by acting a regulator of immune response in skin: promotes Th1/Th17-dominated immune environment contributing to the development of basaloid skin tumors (By similarity). May act as an autoantigen in the immunopathogenesis of psoriasis, with certain peptide regions being a major target for autoreactive T-cells and hence causing their proliferation.

**Cellular Location**

Cytoplasm {ECO:0000250|UniProtKB:Q9QWL7}.

**Tissue Location**

Expressed in the outer root sheath and medulla region of hair follicle specifically from eyebrow and beard, digital pulp, nail matrix and nail bed epithelium, mucosal stratified squamous epithelia and in basal cells of oral epithelium, palmoplantar epidermis and sweat and mammary glands. Also expressed in myoepithelium of prostate, basal layer of urinary bladder, cambial cells of sebaceous gland and in exocervix (at protein level)


**Volume**

125 µl


**KRT17 / CK17 / Cytokeratin 17 Antibody (C-Terminus, clone V21-R) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**KRT17 / CK17 / Cytokeratin 17 Antibody (C-Terminus, clone V21-R) - Images**

Anti-KRT17 / Cytokeratin 17 antibody IHC of human lung.



Anti-KRT17 / Cytokeratin 17 antibody IHC of human prostate.

#### **KRT17 / CK17 / Cytokeratin 17 Antibody (C-Terminus, clone V21-R) - Background**

Type I keratin involved in the formation and maintenance of various skin appendages, specifically in determining shape and orientation of hair (By similarity). Required for the correct growth of hair follicles, in particular for the persistence of the anagen (growth) state (By similarity). Modulates the function of TNF-alpha in the specific context of hair cycling. Regulates protein synthesis and epithelial cell growth through binding to the adapter protein SFN and by stimulating Akt/mTOR pathway (By similarity). Involved in tissue repair. May be a marker of basal cell differentiation in complex epithelia and therefore indicative of a certain type of epithelial "stem cells". Acts as a promoter of epithelial proliferation by acting a regulator of immune response in skin: promotes Th1/Th17-dominated immune environment contributing to the development of basaloid skin tumors (By similarity). May act as an autoantigen in the immunopathogenesis of psoriasis, with certain peptide regions being a major target for autoreactive T-cells and hence causing their proliferation.

#### **KRT17 / CK17 / Cytokeratin 17 Antibody (C-Terminus, clone V21-R) - References**

- Troyanovsky S.M.,et al.Eur. J. Cell Biol. 59:127-137(1992).  
Flohr T.,et al.Eur. J. Immunol. 22:975-979(1992).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Zody M.C.,et al.Nature 440:1045-1049(2006).  
McLean W.H.I.,et al.Nat. Genet. 9:273-278(1995).