

FOXA2 Antibody (C-Terminus) Goat Polyclonal Antibody Catalog # ALS13073

### **Specification**

# FOXA2 Antibody (C-Terminus) - Product Information

Application Primary Accession Reactivity

Host Clonality Calculated MW <u>O9Y261</u> Human, Mouse, Rat, Rabbit, Hamster, Monkey, Pig, Chicken, Horse, Xenopus, Bovine, Dog Goat Polyclonal 48kDa KDa

# FOXA2 Antibody (C-Terminus) - Additional Information

Gene ID 3170

**Other Names** Hepatocyte nuclear factor 3-beta, HNF-3-beta, HNF-3B, Forkhead box protein A2, Transcription factor 3B, TCF-3B, FOXA2, HNF3B, TCF3B

IHC

**Target/Specificity** Human FOXA2. This antibody is expected to recognise both reported isoforms.

**Reconstitution & Storage** Store at -20°C. Minimize freezing and thawing.

**Precautions** FOXA2 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

### **FOXA2 Antibody (C-Terminus) - Protein Information**

Name FOXA2

Synonyms HNF3B, TCF3B

Function

Transcription factor that is involved in embryonic development, establishment of tissue-specific gene expression and regulation of gene expression in differentiated tissues. Is thought to act as a 'pioneer' factor opening the compacted chromatin for other proteins through interactions with nucleosomal core histones and thereby replacing linker histones at target enhancer and/or promoter sites. Binds DNA with the consensus sequence 5'- [AC]A[AT]T[AG]TT[GT][AG][CT]T[CT]-3' (By similarity). In embryonic development is required for notochord formation. Involved in the development of multiple endoderm-derived organ systems such as the liver, pancreas and lungs; FOXA1 and FOXA2 seem to have at least in part redundant roles. Originally described as a



transcription activator for a number of liver genes such as AFP, albumin, tyrosine aminotransferase, PEPCK, etc. Interacts with the cis-acting regulatory regions of these genes. Involved in glucose homeostasis; regulates the expression of genes important for glucose sensing in pancreatic beta- cells and glucose homeostasis. Involved in regulation of fat metabolism. Binds to fibrinogen beta promoter and is involved in IL6- induced fibrinogen beta transcriptional activation.

#### **Cellular Location**


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00089, ECO:0000269|PubMed:14500912}. Cytoplasm Note=Shuttles between the nucleus and cytoplasm in a CRM1-dependent manner; in response to insulin signaling via AKT1 is exported from the nucleus

# FOXA2 Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

# FOXA2 Antibody (C-Terminus) - Images



Anti-FOXA2 antibody IHC of human prostate. FOXA2 Antibody (C-Terminus) - Background

Transcription factor that is involved in embryonic development, establishment of tissue-specific gene expression and regulation of gene expression in differentiated tissues. Is thought to act as a 'pioneer' factor opening the compacted chromatin for other proteins through interactions with nucleosomal core histones and thereby replacing linker histones at target enhancer and/or promoter sites. Binds DNA with the consensus sequence 5'-[AC]A[AT]T[AG]TT[GT][AG][CT]T[CT]-3' (By similarity). In embryonic development is required for notochord formation. Involved in the development of multiple endoderm-derived organ systems such as the liver, pancreas and lungs; FOXA1 and FOXA2 seem to have at least in part redundant roles. Originally described as a transcription activator for a number of liver genes such as AFP, albumin, tyrosine aminotransferase, PEPCK, etc. Interacts with the cis-acting regulatory regions of these genes. Involved in glucose



homeostasis; regulates the expression of genes important for glucose sensing in pancreatic beta-cells and glucose homeostasis. Involved in regulation of fat metabolism. Binds to fibrinogen beta promoter and is involved in IL6-induced fibrinogen beta transcriptional activation.

### FOXA2 Antibody (C-Terminus) - References

Yamada S.,et al.Diabetologia 43:121-124(2000). Hinokio Y.,et al.Submitted (MAY-1999) to the EMBL/GenBank/DDBJ databases. Navas M.A.,et al.Hum. Hered. 50:370-381(2000). Deloukas P.,et al.Nature 414:865-871(2001). Wolfrum C.,et al.Proc. Natl. Acad. Sci. U.S.A. 100:11624-11629(2003).