

**DNMT / DNMT1 Antibody (Internal)**  
**Goat Polyclonal Antibody**  
**Catalog # ALS13338****Specification**

---

**DNMT / DNMT1 Antibody (Internal) - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">P26358</a>
Reactivity	Human, Mouse, Rat, Rabbit, Monkey, Pig, Sheep, Horse, Bovine, Dog
Host	Goat
Clonality	Polyclonal
Calculated MW	183kDa KDa

**DNMT / DNMT1 Antibody (Internal) - Additional Information****Gene ID** 1786**Other Names**

DNA (cytosine-5)-methyltransferase 1, Dnmt1, 2.1.1.37, CXXC-type zinc finger protein 9, DNA methyltransferase Hsa1, DNA MTase Hsa1, M.Hsa1, MCMT, DNMT1, AIM, CXXC9, DNMT

**Target/Specificity**

Human DNMT1.

**Reconstitution & Storage**

Store at -20°C. Minimize freezing and thawing.

**Precautions**

DNMT / DNMT1 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

**DNMT / DNMT1 Antibody (Internal) - Protein Information****Name** DNMT1**Synonyms** AIM, CXXC9, DNMT**Function**

Methylates CpG residues. Preferentially methylates hemimethylated DNA. Associates with DNA replication sites in S phase maintaining the methylation pattern in the newly synthesized strand, that is essential for epigenetic inheritance. Associates with chromatin during G2 and M phases to maintain DNA methylation independently of replication. It is responsible for maintaining methylation patterns established in development. DNA methylation is coordinated with methylation of histones. Mediates transcriptional repression by direct binding to HDAC2. In association with DNMT3B and via the recruitment of CTCFL/BORIS, involved in activation of BAG1 gene expression by modulating dimethylation of promoter histone H3 at H3K4 and H3K9. Probably forms a corepressor complex required for activated KRAS- mediated promoter hypermethylation

and transcriptional silencing of tumor suppressor genes (TSGs) or other tumor-related genes in colorectal cancer (CRC) cells (PubMed:<a href="http://www.uniprot.org/citations/24623306" target="\_blank">24623306</a>). Also required to maintain a transcriptionally repressive state of genes in undifferentiated embryonic stem cells (ESCs) (PubMed:<a href="http://www.uniprot.org/citations/24623306" target="\_blank">24623306</a>). Associates at promoter regions of tumor suppressor genes (TSGs) leading to their gene silencing (PubMed:<a href="http://www.uniprot.org/citations/24623306" target="\_blank">24623306</a>). Promotes tumor growth (PubMed:<a href="http://www.uniprot.org/citations/24623306" target="\_blank">24623306</a>).

**Cellular Location**

Nucleus. Note=Localized to the perinucleolar region.

**Tissue Location**

Ubiquitous; highly expressed in fetal tissues, heart, kidney, placenta, peripheral blood mononuclear cells, and expressed at lower levels in spleen, lung, brain, small intestine, colon, liver, and skeletal muscle. Isoform 2 is less expressed than isoform 1.


**DNMT / DNMT1 Antibody (Internal) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**DNMT / DNMT1 Antibody (Internal) - Images**

Antibody (0.5 ug/ml) staining of Jurkat cell lysate (35 ug protein in RIPA buffer).



Anti-DNMT1 antibody IHC of human kidney.

#### **DNMT / DNMT1 Antibody (Internal) - Background**

Methylates CpG residues. Preferentially methylates hemimethylated DNA. Associates with DNA replication sites in S phase maintaining the methylation pattern in the newly synthesized strand, that is essential for epigenetic inheritance. Associates with chromatin during G2 and M phases to maintain DNA methylation independently of replication. It is responsible for maintaining methylation patterns established in development. DNA methylation is coordinated with methylation of histones. Mediates transcriptional repression by direct binding to HDAC2. In association with DNMT3B and via the recruitment of CTCFL/BORIS, involved in activation of BAG1 gene expression by modulating dimethylation of promoter histone H3 at H3K4 and H3K9.

#### **DNMT / DNMT1 Antibody (Internal) - References**

- Yen R.-W.C., et al. Nucleic Acids Res. 20:2287-2291(1992).  
Yoder J.A., et al. J. Biol. Chem. 271:31092-31097(1996).  
Li L.C., et al. Submitted (AUG-1999) to the EMBL/GenBank/DBJ databases.  
Grimwood J., et al. Nature 428:529-535(2004).  
Hsu D.-W., et al. Proc. Natl. Acad. Sci. U.S.A. 96:9751-9756(1999).