

CD32B Antibody (C-Terminus)
Goat Polyclonal Antibody
Catalog # ALS13425**Specification**

CD32B Antibody (C-Terminus) - Product Information

Application	WB, IHC
Primary Accession	P31994
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Calculated MW	34kDa KDa

CD32B Antibody (C-Terminus) - Additional Information**Gene ID** 2213**Other Names**

Low affinity immunoglobulin gamma Fc region receptor II-b, IgG Fc receptor II-b, CDw32, Fc-gamma RII-b, Fc-gamma-RIIb, FcRII-b, CD32, FCGR2B, CD32, FCG2, IGFR2

Target/Specificity

Human FCGR2B / CD32B. This antibody is expected to recognise all reported isoforms (NP_001002273.1; NP_001002274.1; NP_001002275.1; NP_003992.3; NP_001177757.1).

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.

Precautions

CD32B Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

CD32B Antibody (C-Terminus) - Protein Information**Name** FCGR2B**Synonyms** CD32, FCG2, IGFR2**Function**

Receptor for the Fc region of complexed or aggregated immunoglobulins gamma. Low affinity receptor. Involved in a variety of effector and regulatory functions such as phagocytosis of immune complexes and modulation of antibody production by B-cells. Binding to this receptor results in down-modulation of previous state of cell activation triggered via antigen receptors on B-cells (BCR), T-cells (TCR) or via another Fc receptor. Isoform IIB1 fails to mediate endocytosis or phagocytosis. Isoform IIB2 does not trigger phagocytosis.

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location

Is the most broadly distributed Fc-gamma-receptor. Expressed in monocyte, neutrophils, macrophages, basophils, eosinophils, Langerhans cells, B-cells, platelets cells and placenta (endothelial cells). Not detected in natural killer cells


CD32B Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD32B Antibody (C-Terminus) - Images

Antibody (0.3 ug/ml) staining of K562 lysate (35 ug protein in RIPA buffer).



Anti-CD32B antibody IHC of human liver.

CD32B Antibody (C-Terminus) - Background

Receptor for the Fc region of complexed or aggregated immunoglobulins gamma. Low affinity

receptor. Involved in a variety of effector and regulatory functions such as phagocytosis of immune complexes and modulation of antibody production by B- cells. Binding to this receptor results in down-modulation of previous state of cell activation triggered via antigen receptors on B-cells (BCR), T-cells (TCR) or via another Fc receptor. Isoform IIB1 fails to mediate endocytosis or phagocytosis. Isoform IIB2 does not trigger phagocytosis.

CD32B Antibody (C-Terminus) - References

Ng S.,et al.Submitted (JAN-1997) to the EMBL/GenBank/DDBJ databases.
Stuart S.G.,et al.EMBO J. 8:3657-3666(1989).
Brooks D.G.,et al.J. Exp. Med. 170:1369-1385(1989).
Engelhardt W.,et al.Eur. J. Immunol. 20:1367-1377(1990).
Ebert L.,et al.Submitted (MAY-2004) to the EMBL/GenBank/DDBJ databases.