

**AHR Antibody (aa2-51)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS13675****Specification**

---

**AHR Antibody (aa2-51) - Product Information**

Application	IF, WB, IHC
Primary Accession	<a href="#">P35869</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	96kDa KDa

**AHR Antibody (aa2-51) - Additional Information****Gene ID** 196**Other Names**

Aryl hydrocarbon receptor, Ah receptor, AhR, Class E basic helix-loop-helix protein 76, bHLHe76, AHR, BHLHE76

**Target/Specificity**

AhR (Ab-36) Antibody detects endogenous levels of total AhR protein.

**Reconstitution & Storage**

Store at -20°C for up to one year.

**Precautions**

AHR Antibody (aa2-51) is for research use only and not for use in diagnostic or therapeutic procedures.

**AHR Antibody (aa2-51) - Protein Information****Name** AHR {ECO:0000303|PubMed:8393992, ECO:0000312|HGNC:HGNC:348}**Function**

Ligand-activated transcription factor that enables cells to adapt to changing conditions by sensing compounds from the environment, diet, microbiome and cellular metabolism, and which plays important roles in development, immunity and cancer (PubMed:[30373764](http://www.uniprot.org/citations/30373764), PubMed:[23275542](http://www.uniprot.org/citations/23275542), PubMed:[7961644](http://www.uniprot.org/citations/7961644), PubMed:[32818467](http://www.uniprot.org/citations/32818467)). Upon ligand binding, translocates into the nucleus, where it heterodimerizes with ARNT and induces transcription by binding to xenobiotic response elements (XRE) (PubMed:[30373764](http://www.uniprot.org/citations/30373764), PubMed:[23275542](http://www.uniprot.org/citations/23275542), PubMed:[7961644](http://www.uniprot.org/citations/7961644)). Regulates a

variety of biological processes, including angiogenesis, hematopoiesis, drug and lipid metabolism, cell motility and immune modulation (PubMed:<a href="http://www.uniprot.org/citations/12213388" target="\_blank">12213388</a>). Xenobiotics can act as ligands: upon xenobiotic- binding, activates the expression of multiple phase I and II xenobiotic chemical metabolizing enzyme genes (such as the CYP1A1 gene) (PubMed:<a href="http://www.uniprot.org/citations/7961644" target="\_blank">7961644</a>). Mediates biochemical and toxic effects of halogenated aromatic hydrocarbons (PubMed:<a href="http://www.uniprot.org/citations/7961644" target="\_blank">7961644</a>, PubMed:<a href="http://www.uniprot.org/citations/34521881" target="\_blank">34521881</a>). Next to xenobiotics, natural ligands derived from plants, microbiota, and endogenous metabolism are potent AHR agonists (PubMed:<a href="http://www.uniprot.org/citations/18076143" target="\_blank">18076143</a>). Tryptophan (Trp) derivatives constitute an important class of endogenous AHR ligands (PubMed:<a href="http://www.uniprot.org/citations/32866000" target="\_blank">32866000</a>, PubMed:<a href="http://www.uniprot.org/citations/32818467" target="\_blank">32818467</a>). Acts as a negative regulator of anti-tumor immunity: indoles and kynurenic acid generated by Trp catabolism act as ligand and activate AHR, thereby promoting AHR-driven cancer cell motility and suppressing adaptive immunity (PubMed:<a href="http://www.uniprot.org/citations/32818467" target="\_blank">32818467</a>). Regulates the circadian clock by inhibiting the basal and circadian expression of the core circadian component PER1 (PubMed:<a href="http://www.uniprot.org/citations/28602820" target="\_blank">28602820</a>). Inhibits PER1 by repressing the CLOCK-BMAL1 heterodimer mediated transcriptional activation of PER1 (PubMed:<a href="http://www.uniprot.org/citations/28602820" target="\_blank">28602820</a>). The heterodimer ARNT:AHR binds to core DNA sequence 5'-TGCGTG-3' within the dioxin response element (DRE) of target gene promoters and activates their transcription (PubMed:<a href="http://www.uniprot.org/citations/28602820" target="\_blank">28602820</a>).

#### **Cellular Location**

Cytoplasm. Nucleus. Note=Initially cytoplasmic; upon binding with ligand and interaction with a HSP90, it translocates to the nucleus.

#### **Tissue Location**

Expressed in all tissues tested including blood, brain, heart, kidney, liver, lung, pancreas and skeletal muscle Expressed in retinal photoreceptors (PubMed:29726989)

#### **Volume**


50 µl


#### **AHR Antibody (aa2-51) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **AHR Antibody (aa2-51) - Images**






Immunofluorescence of HeLa cells, using AhR (Ab-36) Antibody.



Western blot of extracts from 293 cells, using AhR (Ab-36) Antibody.



Anti-Aryl Hydrocarbon Receptor antibody IHC of human skin.

### **AHR Antibody (aa2-51) - Background**

Ligand-activated transcriptional activator. Binds to the XRE promoter region of genes it activates. Activates the expression of multiple phase I and II xenobiotic chemical metabolizing enzyme genes (such as the CYP1A1 gene). Mediates biochemical and toxic effects of halogenated aromatic hydrocarbons. Involved in cell-cycle regulation. Likely to play an important role in the development and maturation of many tissues. Regulates the circadian clock by inhibiting the basal and circadian expression of the core circadian component PER1. Inhibits PER1 by repressing the CLOCK-ARNTL/BMAL1 heterodimer mediated transcriptional activation of PER1.

### **AHR Antibody (aa2-51) - References**

- Itoh S.,et al.Nucleic Acids Res. 21:3578-3578(1993).
- Dolwick K.M.,et al.Mol. Pharmacol. 44:911-917(1993).
- Ema M.,et al.J. Biochem. 116:845-851(1994).
- Scherer S.W.,et al.Science 300:767-772(2003).
- Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.