

MMP13 Antibody
Rabbit Polyclonal Antibody
Catalog # ALS13890**Specification**

MMP13 Antibody - Product Information

Application	IHC
Primary Accession	P45452
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54kDa KDa

MMP13 Antibody - Additional Information**Gene ID** 4322**Other Names**

Collagenase 3, 3.4.24.-, Matrix metalloproteinase-13, MMP-13, MMP13

Reconstitution & Storage

Stable for 24 months when stored at 2-8°C.

Precautions

MMP13 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

MMP13 Antibody - Protein Information**Name** MMP13**Function**

Plays a role in the degradation of extracellular matrix proteins including fibrillar collagen, fibronectin, TNC and ACAN. Cleaves triple helical collagens, including type I, type II and type III collagen, but has the highest activity with soluble type II collagen. Can also degrade collagen type IV, type XIV and type X. May also function by activating or degrading key regulatory proteins, such as TGFB1 and CCN2. Plays a role in wound healing, tissue remodeling, cartilage degradation, bone development, bone mineralization and ossification. Required for normal embryonic bone development and ossification. Plays a role in the healing of bone fractures via endochondral ossification. Plays a role in wound healing, probably by a mechanism that involves proteolytic activation of TGFB1 and degradation of CCN2. Plays a role in keratinocyte migration during wound healing. May play a role in cell migration and in tumor cell invasion.

Cellular Location

Secreted, extracellular space, extracellular matrix. Secreted

Tissue Location

Detected in fetal cartilage and calvaria, in chondrocytes of hypertrophic cartilage in vertebrae and in the dorsal end of ribs undergoing ossification, as well as in osteoblasts and periosteal cells

below the inner periosteal region of ossified ribs Detected in chondrocytes from in joint cartilage that have been treated with TNF and IL1B, but not in untreated chondrocytes. Detected in T lymphocytes. Detected in breast carcinoma tissue


Volume
125 µl

MMP13 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MMP13 Antibody - Images



Anti-MMP13 antibody IHC of human prostate.

MMP13 Antibody - Background

Plays a role in the degradation of extracellular matrix proteins including fibrillar collagen, fibronectin, TNC and ACAN. Cleaves triple helical collagens, including type I, type II and type III collagen, but has the highest activity with soluble type II collagen. Can also degrade collagen type IV, type XIV and type X. May also function by activating or degrading key regulatory proteins, such as TGFB1 and CTGF. Plays a role in wound healing, tissue remodeling, cartilage degradation, bone development, bone mineralization and ossification. Required for normal embryonic bone development and ossification. Plays a role in the healing of bone fractures via endochondral ossification. Plays a role in wound healing, probably by a mechanism that involves proteolytic activation of TGFB1 and degradation of CTGF. Plays a role in keratinocyte migration during wound healing. May play a role in cell migration and in tumor cell invasion.

MMP13 Antibody - References

Freije J.M.P., et al. J. Biol. Chem. 269:16766-16773(1994).
Willmroth F., et al. Immunobiology 198:375-384(1998).

Ota T.,et al.Nat. Genet. 36:40-45(2004).
Knaeuper V.,et al.J. Biol. Chem. 271:1544-1550(1996).
Knaeuper V.,et al.J. Biol. Chem. 271:17124-17131(1996).