10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

HDAC6 Antibody (aa7-56)
Rabbit Polyclonal Antibody
Catalog # ALS13928

Specification

HDAC6 Antibody (aa7-56) - Product Information

Application	**IF, WB, IHC**
Primary Accession	Q9UBN7
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**131kDa KDa**

HDAC6 Antibody (aa7-56) - Additional Information

Gene ID 10013

Other Names
Histone deacetylase 6, HD6, 3.5.1.98, HDAC6, KIAA0901

Target/Specificity
HDAC6 (Ab-22) Antibody detects endogenous levels of total HDAC6 protein.

Reconstitution & Storage
Store at -20°C for up to one year.

Precautions
HDAC6 Antibody (aa7-56) is for research use only and not for use in diagnostic or therapeutic procedures.

HDAC6 Antibody (aa7-56) - Protein Information

Name HDAC6 {ECO:0000303|PubMed:10220385, ECO:0000312|HGNC:HGNC:14064}

Function
Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4) (PubMed:10220385). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events (PubMed:10220385). Histone deacetylases act via the formation of large multiprotein complexes (PubMed:10220385). In addition to histones, deacetylates other proteins: plays a central role in microtubule-dependent cell motility by mediating deacetylation of tubulin (PubMed:12024216, PubMed:20308065, PubMed:26246421). Required for cilia disassembly; via deacetylation of alpha-tubulin (PubMed:<a

href="http://www.uniprot.org/citations/17604723" target="_blank">17604723, PubMed:26246421). Promotes deacetylation of CTTN, leading to actin polymerization, promotion of autophagosome-lysosome fusion and completion of autophagy (PubMed:30538141). Involved in the MTA1-mediated epigenetic regulation of ESR1 expression in breast cancer (PubMed:24413532). Promotes odontoblast differentiation following IPO7-mediated nuclear import and subsequent repression of RUNX2 expression (By similarity). In addition to its protein deacetylase activity, plays a key role in the degradation of misfolded proteins: when misfolded proteins are too abundant to be degraded by the chaperone refolding system and the ubiquitin-proteasome, mediates the transport of misfolded proteins to a cytoplasmic juxtanuclear structure called aggresome (PubMed:17846173). Probably acts as an adapter that recognizes polyubiquitinated misfolded proteins and target them to the aggresome, facilitating their clearance by autophagy (PubMed:17846173).

Cellular Location
Cytoplasm. Cytoplasm, cytoskeleton. Nucleus {ECO:0000250|UniProtKB:Q9Z2V5}. Perikaryon {ECO:0000250|UniProtKB:Q9Z2V5}. Cell projection, dendrite {ECO:0000250|UniProtKB:Q9Z2V5}. Cell projection, axon {ECO:0000250|UniProtKB:Q9Z2V5}. Cell projection, cilium. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, cilium basal body. Note=It is mainly cytoplasmic, where it is associated with microtubules


Volume
50 µl

HDAC6 Antibody (aa7-56) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

HDAC6 Antibody (aa7-56) - Images




Immunofluorescence of HepG2 cells, using HDAC6 (Ab-22) Antibody.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Western blot of extracts from HepG2 cells, using HDAC6 (Ab-22) Antibody.



Anti-HDAC6 antibody IHC of human intestine, ganglion cells.

HDAC6 Antibody (aa7-56) - Background

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes (By similarity). Plays a central role in microtubule-dependent cell motility via deacetylation of tubulin. Involved in the MTA1-mediated epigenetic regulation of ESR1 expression in breast cancer.

HDAC6 Antibody (aa7-56) - References

Grozinger C.M.,et al.Proc. Natl. Acad. Sci. U.S.A. 96:4868-4873(1999).
Nagase T.,et al.DNA Res. 5:355-364(1998).
Ohara O.,et al.Submitted (JAN-2004) to the EMBL/GenBank/DDBJ databases.
Strom T.M.,et al.Submitted (OCT-1998) to the EMBL/GenBank/DDBJ databases.
Ross M.T.,et al.Nature 434:325-337(2005).