

WDR92 Antibody (Internal)
Rabbit Polyclonal Antibody
Catalog # ALS13979**Specification**

WDR92 Antibody (Internal) - Product Information

Application	IF, WB
Primary Accession	Q96MX6
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40kDa KDa

WDR92 Antibody (Internal) - Additional Information**Gene ID** 116143**Other Names**

WD repeat-containing protein 92, WD repeat-containing protein Monad, WDR92

Target/Specificity

Human WDR92

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

Precautions

WDR92 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

WDR92 Antibody (Internal) - Protein Information**Name** DNAAF10 ([HGNC:25176](#))**Synonyms** WDR92**Function**

Key assembly factor specifically required for the stability of axonemal dynein heavy chains in cytoplasm.

Cellular Location

Dynein axonemal particle {ECO:0000250|UniProtKB:A8J3F6}

Tissue Location


Widely expressed with the highest expression in testis.

WDR92 Antibody (Internal) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

WDR92 Antibody (Internal) - Images



Immunofluorescence of WDR92 in human testis tissue with WDR92 antibody at 20 ug/ml.



Western blot analysis of WDR92 in human kidney tissue lysate with WDR92 antibody at (A) 1 and...

WDR92 Antibody (Internal) - Background

Seems to act as a modulator of apoptosis.

WDR92 Antibody (Internal) - References

Ota T.,et al.Nat. Genet. 36:40-45(2004).
Hillier L.W.,et al.Nature 434:724-731(2005).
Saeki M.,et al.Biochem. Biophys. Res. Commun. 342:568-572(2006).
Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).
Xu C.,et al.Protein Cell 2:202-214(2011).