

USP28 Antibody (Internal)
Goat Polyclonal Antibody
Catalog # ALS14572**Specification**

USP28 Antibody (Internal) - Product Information

Application	IHC
Primary Accession	Q96RU2
Reactivity	Human, Rabbit, Monkey, Pig, Horse, Bovine, Dog
Host	Goat
Clonality	Polyclonal
Calculated MW	122kDa KDa

USP28 Antibody (Internal) - Additional Information**Gene ID** 57646**Other Names**

Ubiquitin carboxyl-terminal hydrolase 28, 3.4.19.12, Deubiquitinating enzyme 28, Ubiquitin thioesterase 28, Ubiquitin-specific-processing protease 28, USP28, KIAA1515

Target/Specificity

Human USP28.

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

USP28 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

USP28 Antibody (Internal) - Protein Information**Name** USP28**Synonyms** KIAA1515**Function**

Deubiquitinase involved in DNA damage response checkpoint and MYC proto-oncogene stability. Involved in DNA damage induced apoptosis by specifically deubiquitinating proteins of the DNA damage pathway such as CLSPN. Also involved in G2 DNA damage checkpoint, by deubiquitinating CLSPN, and preventing its degradation by the anaphase promoting complex/cyclosome (APC/C). In contrast, it does not deubiquitinate PLK1. Specifically deubiquitinates MYC in the nucleoplasm, leading to prevent MYC degradation by the proteasome: acts by specifically interacting with isoform 1 of FBXW7 (FBW7alpha) in the nucleoplasm and counteracting ubiquitination of MYC by the SCF(FBW7) complex. In contrast, it does not interact with isoform 4 of FBXW7 (FBW7gamma) in the nucleolus, allowing MYC degradation and explaining the selective MYC degradation in the

nucleolus. Deubiquitinates ZNF304, hence preventing ZNF304 degradation by the proteasome and leading to the activated KRAS-mediated promoter hypermethylation and transcriptional silencing of tumor suppressor genes (TSGs) in a subset of colorectal cancers (CRC) cells (PubMed:24623306).

Cellular Location


Nucleus, nucleoplasm

USP28 Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

USP28 Antibody (Internal) - Images



Anti-USP28 antibody IHC of human uterus.

USP28 Antibody (Internal) - Background

Deubiquitinase involved in DNA damage response checkpoint and MYC proto-oncogene stability. Involved in DNA damage induced apoptosis by specifically deubiquitinating proteins of the DNA damage pathway such as CLSPN. Also involved in G2 DNA damage checkpoint, by deubiquitinating CLSPN, and preventing its degradation by the anaphase promoting complex/cyclosome (APC/C). In contrast, it does not deubiquitinate PLK1. Specifically deubiquitinates MYC in the nucleoplasm, leading to prevent MYC degradation by the proteasome: acts by specifically interacting with isoform 1 of FBXW7 (FBW7alpha) in the nucleoplasm and counteracting ubiquitination of MYC by the SCF(FBW7) complex. In contrast, it does not interact with isoform 4 of FBXW7 (FBW7gamma) in the nucleolus, allowing MYC degradation and explaining the selective MYC degradation in the nucleolus.

USP28 Antibody (Internal) - References

Valero R., et al. Genome Biol. 2:RESEARCH0043.1-RESEARCH0043.10(2001).

Taylor T.D.,et al.Nature 440:497-500(2006).

Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

Nagase T.,et al.DNA Res. 7:143-150(2000).

Zhang D.,et al.Cell 126:529-542(2006).