

FN1 / Fibronectin Antibody (clone 23C9)
Mouse Monoclonal Antibody
Catalog # ALS14803**Specification**

FN1 / Fibronectin Antibody (clone 23C9) - Product Information

Application	WB
Primary Accession	P02751
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	263kDa KDa

FN1 / Fibronectin Antibody (clone 23C9) - Additional Information**Gene ID** 2335**Other Names**

Fibronectin, FN, Cold-insoluble globulin, CIG, Anastellin, Ugl-Y1, Ugl-Y2, Ugl-Y3, FN1, FN

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.

Precautions

FN1 / Fibronectin Antibody (clone 23C9) is for research use only and not for use in diagnostic or therapeutic procedures.

FN1 / Fibronectin Antibody (clone 23C9) - Protein Information**Name** FN1 ([HGNC:3778](#))**Synonyms** FN**Function**

Fibronectins bind cell surfaces and various compounds including collagen, fibrin, heparin, DNA, and actin (PubMed: [3024962](http://www.uniprot.org/citations/3024962), PubMed: [3900070](http://www.uniprot.org/citations/3900070), PubMed: [3593230](http://www.uniprot.org/citations/3593230), PubMed: [7989369](http://www.uniprot.org/citations/7989369)). Fibronectins are involved in cell adhesion, cell motility, opsonization, wound healing, and maintenance of cell shape (PubMed: [3024962](http://www.uniprot.org/citations/3024962), PubMed: [3900070](http://www.uniprot.org/citations/3900070), PubMed: [3593230](http://www.uniprot.org/citations/3593230), PubMed: [7989369](http://www.uniprot.org/citations/7989369)). Involved in osteoblast compaction through the fibronectin fibrillogenesis cell-mediated matrix assembly process, essential for osteoblast mineralization (By similarity). Participates in the regulation of type I collagen deposition by osteoblasts (By similarity). Acts as a ligand for the LILRB4 receptor,

inhibiting FCGR1A/CD64-mediated monocyte activation (PubMed:34089617).

Cellular Location

Secreted, extracellular space, extracellular matrix. Secreted {ECO:0000250|UniProtKB:P11276}

Tissue Location

Expressed in the inner limiting membrane and around blood vessels in the retina (at protein level) (PubMed:29777959) Plasma FN (soluble dimeric form) is secreted by hepatocytes. Cellular FN (dimeric or cross-linked multimeric forms), made by fibroblasts, epithelial and other cell types, is deposited as fibrils in the extracellular matrix. Ugl-Y1, Ugl-Y2 and Ugl-Y3 are found in urine (PubMed:17614963).

Volume


50 µl

FN1 / Fibronectin Antibody (clone 23C9) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FN1 / Fibronectin Antibody (clone 23C9) - Images



0

FN1 / Fibronectin Antibody (clone 23C9) - Background

Fibronectins bind cell surfaces and various compounds including collagen, fibrin, heparin, DNA, and actin. Fibronectins are involved in cell adhesion, cell motility, opsonization, wound healing, and maintenance of cell shape. Involved in osteoblast compaction through the fibronectin fibrillogenesis cell-mediated matrix assembly process, essential for osteoblast mineralization. Participates in the regulation of type I collagen deposition by osteoblasts.

FN1 / Fibronectin Antibody (clone 23C9) - References

Schor S.L.,et al.Breast Cancer Res. 3:373-379(2001).

Kay R.A.,et al.Cancer Res. 65:10742-10749(2005).

Kato S.,et al.DNA Res. 12:53-62(2005).

Totoki Y.,et al.Submitted (MAR-2005) to the EMBL/GenBank/DDBJ databases.

Bechtel S.,et al.BMC Genomics 8:399-399(2007).