

HAMP / Hepcidin Antibody (N-Terminus)
Goat Polyclonal Antibody
Catalog # ALS14957**Specification**

HAMP / Hepcidin Antibody (N-Terminus) - Product Information

Application	IHC
Primary Accession	P81172
Reactivity	Human, Monkey
Host	Goat
Clonality	Polyclonal
Calculated MW	9kDa KDa

HAMP / Hepcidin Antibody (N-Terminus) - Additional Information**Gene ID** 57817**Other Names**

Hepcidin, Liver-expressed antimicrobial peptide 1, LEAP-1, Putative liver tumor regressor, PLTR, Hepcidin-25, Hepc25, Hepcidin-20, Hepc20, HAMP, HEPC, LEAP1

Target/Specificity

Human HAMP / Hepcidin.

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

HAMP / Hepcidin Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

HAMP / Hepcidin Antibody (N-Terminus) - Protein Information**Name** HAMP ([HGNC:15598](#))**Synonyms** HEPC, LEAP1**Function**

Liver-produced hormone that constitutes the main circulating regulator of iron absorption and distribution across tissues. Acts by promoting endocytosis and degradation of ferroportin/SLC40A1, leading to the retention of iron in iron-exporting cells and decreased flow of iron into plasma (PubMed:22682227, PubMed:29237594, PubMed:32814342). Controls the major flows of iron into plasma: absorption of dietary iron in the intestine, recycling of iron by macrophages, which phagocytose old erythrocytes and other cells, and mobilization of stored iron from hepatocytes (PubMed:22306005).


Cellular Location
Secreted.**Tissue Location**

Highest expression in liver and to a lesser extent in heart and brain. Low levels in lung, tonsils, salivary gland, trachea, prostate gland, adrenal gland and thyroid gland. Secreted into the urine and blood (PubMed:11034317). Expressed by hepatocytes (PubMed:15124018).

HAMP / Hepcidin Antibody (N-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HAMP / Hepcidin Antibody (N-Terminus) - Images

Anti-HAMP / Hepcidin antibody IHC of human heart.

HAMP / Hepcidin Antibody (N-Terminus) - Background

Liver-produced hormone that constitutes the main circulating regulator of iron absorption and distribution across tissues. Acts by promoting endocytosis and degradation of ferroportin, leading to the retention of iron in iron-exporting cells and decreased flow of iron into plasma. Controls the major flows of iron into plasma: absorption of dietary iron in the intestine, recycling of iron by macrophages, which phagocytose old erythrocytes and other cells, and mobilization of stored iron from hepatocytes (PubMed:22306005).

HAMP / Hepcidin Antibody (N-Terminus) - References

Park C.H., et al. J. Biol. Chem. 276:7806-7810(2001).
Krause A., et al. FEBS Lett. 480:147-150(2000).
Jung J.-W., et al. Submitted (FEB-1999) to the EMBL/GenBank/DBJ databases.
Clark H.F., et al. Genome Res. 13:2265-2270(2003).
Grimwood J., et al. Nature 428:529-535(2004).

