

#### GJA1 / CX43 / Connexin 43 Antibody (aa333-382) Rabbit Polyclonal Antibody Catalog # ALS15038

### Specification

# GJA1 / CX43 / Connexin 43 Antibody (aa333-382) - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW IF, IHC <u>P17302</u> Human, Mouse, Rat Rabbit Polyclonal 43kDa KDa

#### GJA1 / CX43 / Connexin 43 Antibody (aa333-382) - Additional Information

Gene ID 2697

Other Names Gap junction alpha-1 protein, Connexin-43, Cx43, Gap junction 43 kDa heart protein, GJA1, GJAL

**Target/Specificity** Connexin 43 Antibody detects endogenous levels of total Connexin 43 protein.

**Reconstitution & Storage** Store at -20°C for up to one year.

**Precautions** GJA1 / CX43 / Connexin 43 Antibody (aa333-382) is for research use only and not for use in diagnostic or therapeutic procedures.

#### GJA1 / CX43 / Connexin 43 Antibody (aa333-382) - Protein Information

Name GJA1

Synonyms GJAL

Function

Gap junction protein that acts as a regulator of bladder capacity. A gap junction consists of a cluster of closely packed pairs of transmembrane channels, the connexons, through which materials of low MW diffuse from one cell to a neighboring cell. May play a critical role in the physiology of hearing by participating in the recycling of potassium to the cochlear endolymph. Negative regulator of bladder functional capacity: acts by enhancing intercellular electrical and chemical transmission, thus sensitizing bladder muscles to cholinergic neural stimuli and causing them to contract (By similarity). May play a role in cell growth inhibition through the regulation of NOV expression and localization. Plays an essential role in gap junction communication in the ventricles (By similarity).

**Cellular Location** 



Cell membrane; Multi-pass membrane protein. Cell junction, gap junction. Endoplasmic reticulum {ECO:0000250|UniProtKB:P23242}. Note=Localizes at the intercalated disk (ICD) in cardiomyocytes and the proper localization at ICD is dependent on TMEM65. {ECO:0000250|UniProtKB:P23242}

**Tissue Location** Expressed in the heart and fetal cochlea.


Volume 50 μl

# GJA1 / CX43 / Connexin 43 Antibody (aa333-382) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

GJA1 / CX43 / Connexin 43 Antibody (aa333-382) - Images



Immunofluorescence of A549 cells, using Connexin 43 Antibody.



Anti-GJA1 / Connexin 43 antibody IHC of human heart. GJA1 / CX43 / Connexin 43 Antibody (aa333-382) - Background

Gap junction protein that acts as a regulator of bladder capacity. A gap junction consists of a cluster of closely packed pairs of transmembrane channels, the connexons, through which materials of low MW diffuse from one cell to a neighboring cell. May play a critical role in the physiology of



hearing by participating in the recycling of potassium to the cochlear endolymph. Negative regulator of bladder functional capacity: acts by enhancing intercellular electrical and chemical transmission, thus sensitizing bladder muscles to cholinergic neural stimuli and causing them to contract (By similarity).

### GJA1 / CX43 / Connexin 43 Antibody (aa333-382) - References

Fishman G.I.,et al.J. Cell Biol. 111:589-598(1990). Fishman G.I.,et al.Genomics 10:250-256(1991). Haefliger J.-A.,et al.Eur. Heart J. 20:1843-1843(1999). Halleck A.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases. Ota T.,et al.Nat. Genet. 36:40-45(2004).