

**PPIF / Cyclophilin F Antibody (aa86-135)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS15105****Specification**

---

**PPIF / Cyclophilin F Antibody (aa86-135) - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">P30405</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22kDa KDa

**PPIF / Cyclophilin F Antibody (aa86-135) - Additional Information****Gene ID** 10105**Other Names**

Peptidyl-prolyl cis-trans isomerase F, mitochondrial, PPlase F, 5.2.1.8, Cyclophilin D, CyP-D, CypD, Cyclophilin F, Mitochondrial cyclophilin, CyP-M, Rotamase F, PPIF, CYP3

**Target/Specificity**

PPIF Antibody detects endogenous levels of total PPIF protein.

**Reconstitution & Storage**

Store at -20°C for up to one year.

**Precautions**

PPIF / Cyclophilin F Antibody (aa86-135) is for research use only and not for use in diagnostic or therapeutic procedures.

**PPIF / Cyclophilin F Antibody (aa86-135) - Protein Information****Name** PPIF**Synonyms** CYP3**Function**

PPlase that catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and may therefore assist protein folding (PubMed:[20676357](http://www.uniprot.org/citations/20676357)). Involved in regulation of the mitochondrial permeability transition pore (mPTP) (PubMed:[26387735](http://www.uniprot.org/citations/26387735)). It is proposed that its association with the mPTP is masking a binding site for inhibiting inorganic phosphate (Pi) and promotes the open probability of the mPTP leading to apoptosis or necrosis; the requirement of the PPlase activity for this function is debated (PubMed:[26387735](http://www.uniprot.org/citations/26387735)). In cooperation with mitochondrial p53/TP53 is involved in activating oxidative stress-induced

necrosis (PubMed:<a href="http://www.uniprot.org/citations/22726440" target="\_blank">22726440</a>). Involved in modulation of mitochondrial membrane F(1)F(0) ATP synthase activity and regulation of mitochondrial matrix adenine nucleotide levels (By similarity). Has anti-apoptotic activity independently of mPTP and in cooperation with BCL2 inhibits cytochrome c-dependent apoptosis (PubMed:<a href="http://www.uniprot.org/citations/19228691" target="\_blank">19228691</a>).

**Cellular Location**

Mitochondrion matrix


**Volume**

50 µl


**PPIF / Cyclophilin F Antibody (aa86-135) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**PPIF / Cyclophilin F Antibody (aa86-135) - Images**

Western blot of extracts from HepG2/HeLa/293 cells, using PPIF Antibody.



Anti-PPIF / Cyclophilin F antibody IHC of human adrenal.

### **PPIF / Cyclophilin F Antibody (aa86-135) - Background**

PPlases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. Involved in regulation of the mitochondrial permeability transition pore (mPTP). It is proposed that its association with the mPTP is masking a binding site for inhibiting inorganic phosphate (Pi) and promotes the open probability of the mPTP leading to apoptosis or necrosis; the requirement of the PPlase activity for this function is debated. In cooperation with mitochondrial TP53 is involved in activating oxidative stress- induced necrosis. Involved in modulation of mitochondrial membrane F(1)F(0) ATP synthase activity and regulation of mitochondrial matrix adenine nucleotide levels. Has anti-apoptotic activity independently of mPTP and in cooperation with BCL2 inhibits cytochrome c-dependent apoptosis.

### **PPIF / Cyclophilin F Antibody (aa86-135) - References**

Bergsma D.J.,et al.J. Biol. Chem. 266:23204-23214(1991).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Deloukas P.,et al.Nature 429:375-381(2004).  
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.  
Johnson N.,et al.Eur. J. Biochem. 263:353-359(1999).