

FNBP1 / FBP17 Antibody (aa30-80)
Rabbit Polyclonal Antibody
Catalog # ALS15245**Specification**

FNBP1 / FBP17 Antibody (aa30-80) - Product Information

Application	WB, IHC
Primary Accession	Q96RU3
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	71kDa KDa

FNBP1 / FBP17 Antibody (aa30-80) - Additional Information**Gene ID** 23048**Other Names**

Formin-binding protein 1, Formin-binding protein 17, hFBP17, FNBP1, FBP17, KIAA0554

Target/Specificity

Human FNBP1

Reconstitution & Storage

Store at 4°C for short term applications. For long term storage, aliquot and store at -20°C.

Precautions

FNBP1 / FBP17 Antibody (aa30-80) is for research use only and not for use in diagnostic or therapeutic procedures.

FNBP1 / FBP17 Antibody (aa30-80) - Protein Information**Name** FNBP1**Synonyms** FBP17, KIAA0554**Function**

May act as a link between RND2 signaling and regulation of the actin cytoskeleton (By similarity). Required to coordinate membrane tubulation with reorganization of the actin cytoskeleton during the late stage of clathrin-mediated endocytosis. Binds to lipids such as phosphatidylinositol 4,5-bisphosphate and phosphatidylserine and promotes membrane invagination and the formation of tubules. Also enhances actin polymerization via the recruitment of WASL/N-WASP, which in turn activates the Arp2/3 complex. Actin polymerization may promote the fission of membrane tubules to form endocytic vesicles. May be required for the lysosomal retention of FASLG/FASL.

Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton. Cytoplasm, cell cortex. Lysosome. Cytoplasmic vesicle. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Membrane, clathrin-coated pit

Note=Enriched in cortical regions coincident with F-actin. Also localizes to endocytic vesicles and lysosomes

Tissue Location


Very highly expressed in the epithelial cells of the gastrointestinal tract, respiratory, reproductive and urinary systems. Also highly expressed in brown adipose tissue, cardiomyocytes, enteric ganglia and glucagon producing cells of the pancreas. Expressed in germ cells of the testis and all regions of the brain

FNBP1 / FBP17 Antibody (aa30-80) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FNBP1 / FBP17 Antibody (aa30-80) - Images



Western blot of FNBP1 in human liver lysate in the 1) absence and 2) presence of immunizing...



Anti-FNBP1 / FBP17 antibody IHC of human brain, cortex neurons.

FNBP1 / FBP17 Antibody (aa30-80) - Background

May act as a link between RND2 signaling and regulation of the actin cytoskeleton (By similarity). Required to coordinate membrane tubulation with reorganization of the actin cytoskeleton during the late stage of clathrin-mediated endocytosis. Binds to lipids such as phosphatidylinositol 4,5-bisphosphate and phosphatidylserine and promotes membrane invagination and the formation of tubules. Also enhances actin polymerization via the recruitment of WASL/N-WASP, which in turn activates the Arp2/3 complex. Actin polymerization may promote the fission of membrane tubules to form endocytic vesicles. May be required for the lysosomal retention of FASLG/FASL.

FNBP1 / FBP17 Antibody (aa30-80) - References

Fuchs U.,et al.Proc. Natl. Acad. Sci. U.S.A. 98:8756-8761(2001).
Nagase T.,et al.DNA Res. 5:31-39(1998).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Humphray S.J.,et al.Nature 429:369-374(2004).
Richnau N.,et al.J. Biol. Chem. 276:35060-35070(2001).