

EPM2A / Laforin Antibody (aa80-130)
Rabbit Polyclonal Antibody
Catalog # ALS15246**Specification**

EPM2A / Laforin Antibody (aa80-130) - Product Information

Application	WB, IHC
Primary Accession	O95278
Reactivity	Human, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	37kDa KDa

EPM2A / Laforin Antibody (aa80-130) - Additional Information**Gene ID** 7957**Other Names**

Laforin, 3.1.3.-, 3.1.3.16, 3.1.3.48, Glucan phosphatase, Lafora PTPase, LAFPTPase, EPM2A

Target/Specificity

Human EPM2A

Reconstitution & Storage

Store at 4°C for short term applications. For long term storage, aliquot and store at -20°C.

Precautions

EPM2A / Laforin Antibody (aa80-130) is for research use only and not for use in diagnostic or therapeutic procedures.

EPM2A / Laforin Antibody (aa80-130) - Protein Information**Name** EPM2A**Function**

Plays an important role in preventing glycogen hyperphosphorylation and the formation of insoluble aggregates, via its activity as glycogen phosphatase, and by promoting the ubiquitination of proteins involved in glycogen metabolism via its interaction with the E3 ubiquitin ligase NHLRC1/malin. Shows strong phosphatase activity towards complex carbohydrates in vitro, avoiding glycogen hyperphosphorylation which is associated with reduced branching and formation of insoluble aggregates (PubMed: [16901901](http://www.uniprot.org/citations/16901901) target="_blank">16901901, PubMed: [23922729](http://www.uniprot.org/citations/23922729) target="_blank">23922729, PubMed: [26231210](http://www.uniprot.org/citations/26231210) target="_blank">26231210, PubMed: [25538239](http://www.uniprot.org/citations/25538239) target="_blank">25538239, PubMed: [25544560](http://www.uniprot.org/citations/25544560) target="_blank">25544560). Dephosphorylates phosphotyrosine and synthetic substrates, such as para- nitrophenylphosphate (pNPP), and has low activity with phosphoserine and phosphothreonine substrates (in vitro) (PubMed: <a

[11001928](http://www.uniprot.org/citations/11001928), PubMed: [11220751](http://www.uniprot.org/citations/11220751), PubMed: [11739371](http://www.uniprot.org/citations/11739371), PubMed: [14532330](http://www.uniprot.org/citations/14532330), PubMed: [16971387](http://www.uniprot.org/citations/16971387), PubMed: [18617530](http://www.uniprot.org/citations/18617530), PubMed: [22036712](http://www.uniprot.org/citations/22036712), PubMed: [23922729](http://www.uniprot.org/citations/23922729), PubMed: [14722920](http://www.uniprot.org/citations/14722920)). Has been shown to dephosphorylate MAPT (By similarity). Forms a complex with NHLRC1/malin and HSP70, which suppresses the cellular toxicity of misfolded proteins by promoting their degradation through the ubiquitin-proteasome system (UPS). Acts as a scaffold protein to facilitate PPP1R3C/PTG ubiquitination by NHLRC1/malin (PubMed: [23922729](http://www.uniprot.org/citations/23922729)). Also promotes proteasome-independent protein degradation through the macroautophagy pathway (PubMed: [20453062](http://www.uniprot.org/citations/20453062)).

Cellular Location

Cytoplasm. Note=Under glycogenolytic conditions localizes to the nucleus [Isoform 2]: Cytoplasm. Endoplasmic reticulum membrane; Peripheral membrane protein; Cytoplasmic side. Cell membrane. Nucleus. Note=Also found in the nucleus. [Isoform 5]: Cytoplasm. Nucleus

Tissue Location


Expressed in heart, skeletal muscle, kidney, pancreas and brain. Isoform 4 is also expressed in the placenta

EPM2A / Laforin Antibody (aa80-130) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

EPM2A / Laforin Antibody (aa80-130) - Images



Western blot of EPM2A in human heart lysate in the 1) absence and 2) presence of immunizing...



Anti-EPM2A / Laforin antibody IHC of human heart.

EPM2A / Laforin Antibody (aa80-130) - Background

Has both dual-specificity protein phosphatase and glucan phosphatase activities. Together with the E3 ubiquitin ligase NHLRC1/malin, appears to be involved in the clearance of toxic polyglucosan and protein aggregates via multiple pathways. Dephosphorylates phosphotyrosine, phosphoserine and phosphothreonine substrates in vitro. Has also been shown to dephosphorylate MAPT. Shows strong phosphatase activity towards complex carbohydrates in vitro, avoiding glycogen hyperphosphorylation which is associated with reduced branching and formation of insoluble aggregates. Forms a complex with NHLRC1/malin and HSP70, which suppresses the cellular toxicity of misfolded proteins by promoting their degradation through the ubiquitin-proteasome system (UPS). Acts as a scaffold protein to facilitate PPP1R3C/PTG ubiquitination by NHLRC1/malin. Also promotes proteasome-independent protein degradation through the macroautophagy pathway. Isoform 2, an inactive phosphatase, could function as a dominant-negative regulator for the phosphatase activity of isoform 1.

EPM2A / Laforin Antibody (aa80-130) - References

Minassian B.A., et al. Nat. Genet. 20:171-174(1998).
Ganesh S., et al. Hum. Mol. Genet. 9:2251-2261(2000).
Lee J.R., et al. Submitted (AUG-1998) to the EMBL/GenBank/DDBJ databases.
Ganesh S., et al. Submitted (NOV-2001) to the EMBL/GenBank/DDBJ databases.
Ota T., et al. Nat. Genet. 36:40-45(2004).