

SS18 Antibody (aa1-50)
Rabbit Polyclonal Antibody
Catalog # ALS15300**Specification**

SS18 Antibody (aa1-50) - Product Information

Application	IHC
Primary Accession	Q15532
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46kDa KDa

SS18 Antibody (aa1-50) - Additional Information**Gene ID** 6760**Other Names**

Protein SSXT, Protein SYT, Synovial sarcoma translocated to X chromosome protein, SS18, SSXT, SYT

Target/Specificity

SSXT Antibody detects endogenous levels of total SSXT protein.

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.

Precautions

SS18 Antibody (aa1-50) is for research use only and not for use in diagnostic or therapeutic procedures.

SS18 Antibody (aa1-50) - Protein Information**Name** SS18**Synonyms** SSXT, SYT**Function**

Appears to function synergistically with RBM14 as a transcriptional coactivator. Isoform 1 and isoform 2 function in nuclear receptor coactivation. Isoform 1 and isoform 2 function in general transcriptional coactivation. Component of SWI/SNF chromatin remodeling subcomplex GBAF that carries out key enzymatic activities, changing chromatin structure by altering DNA-histone contacts within a nucleosome in an ATP-dependent manner (PubMed:29374058).

Cellular Location

Nucleus.

Tissue Location

Fairly ubiquitously expressed. Expressed in synovial sarcomas and in other human cell lines. The fusion genes SSXT- SSX1 and SSXT-SSX2 are expressed only in synovial sarcomas


Volume

50 µl

SS18 Antibody (aa1-50) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SS18 Antibody (aa1-50) - Images

Anti-SS18 / SYT antibody IHC of human prostate.

SS18 Antibody (aa1-50) - Background

Appears to function synergistically with RBM14 as a transcriptional coactivator. Isoform 1 and isoform 2 function in nuclear receptor coactivation. Isoform 1 and isoform 2 function in general transcriptional coactivation.

SS18 Antibody (aa1-50) - References

Clark J., et al. Nat. Genet. 7:502-508(1994).
Brodin B., et al. Gene 268:173-182(2001).
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Fligman I., et al. Am. J. Pathol. 147:1592-1599(1995).
de Bruijn D.R., et al. Oncogene 20:3281-3289(2001).