

GRIK1 / GLUR5 Antibody (Internal) Goat Polyclonal Antibody

Catalog # ALS15357

Specification

GRIK1 / GLUR5 Antibody (Internal) - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW WB, IHC <u>P39086</u> Human, Mouse, Rat, Monkey, Horse Goat Polyclonal 104kDa KDa

GRIK1 / GLUR5 Antibody (Internal) - Additional Information

Gene ID 2897

Other Names Glutamate receptor ionotropic, kainate 1, GluK1, Excitatory amino acid receptor 3, EAA3, Glutamate receptor 5, GluR-5, GluR5, GRIK1, GLUR5

Target/Specificity Human GRIK1. This antibody is expected to recognise both reported isoforms according to NP_000821.1 and NP_783300.1.

Reconstitution & Storage Store at -20°C. Minimize freezing and thawing.

Precautions GRIK1 / GLUR5 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

GRIK1 / GLUR5 Antibody (Internal) - Protein Information

Name GRIK1

Synonyms GLUR5

Function

lonotropic glutamate receptor. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist. May be involved in the transmission of light information from the retina to the hypothalamus.

Cellular Location

Cell membrane; Multi-pass membrane protein. Postsynaptic cell membrane; Multi-pass membrane



protein

Tissue Location

Most abundant in the cerebellum and the suprachiasmatic nuclei (SCN) of the hypothalamus

GRIK1 / GLUR5 Antibody (Internal) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

GRIK1 / GLUR5 Antibody (Internal) - Images



Antibody (1 ug/ml) staining of Human Brain (Cerebellum) lysate (35 ug protein in RIPA buffer).



Anti-GRIK1 / GLUR5 antibody IHC of human brain, cortex.

GRIK1 / GLUR5 Antibody (Internal) - Background

lonotropic glutamate receptor. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L- glutamate



induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist. May be involved in the transmission of light information from the retina to the hypothalamus.

GRIK1 / GLUR5 Antibody (Internal) - References

Gregor P., et al.NeuroReport 4:1343-1346(1993). Korczak B., et al.Recept. Channels 3:41-49(1995). Langer A., et al.Submitted (SEP-1999) to the EMBL/GenBank/DDBJ databases. Nutt S.L., et al.NeuroReport 5:2625-2629(1994). Shibata H., et al.Psychiatr. Genet. 11:139-144(2001).