

UBASH3A / CLIP4 Antibody (Internal)
Goat Polyclonal Antibody
Catalog # ALS15401**Specification**

UBASH3A / CLIP4 Antibody (Internal) - Product Information

Application	WB, IHC
Primary Accession	P57075
Reactivity	Human, Monkey
Host	Goat
Clonality	Polyclonal
Calculated MW	74kDa KDa

UBASH3A / CLIP4 Antibody (Internal) - Additional Information**Gene ID** 53347**Other Names**

Ubiquitin-associated and SH3 domain-containing protein A, Cbl-interacting protein 4, CLIP4, Suppressor of T-cell receptor signaling 2, STS-2, T-cell ubiquitin ligand 1, TULA-1, UBASH3A, STS2

Target/Specificity

Human UBASH3A. This antibody is expected to recognize both reported isoforms (NP_061834.1; NP_001001895.1).

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

UBASH3A / CLIP4 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

UBASH3A / CLIP4 Antibody (Internal) - Protein Information**Name** UBASH3A**Synonyms** STS2**Function**

Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors, EGFR and PDGFRB, on the cell surface. Exhibits negligible protein tyrosine phosphatase activity at neutral pH. May act as a dominant-negative regulator of UBASH3B- dependent dephosphorylation. May inhibit dynamin-dependent endocytic pathways by functionally sequestering dynamin via its SH3 domain.

Cellular Location

Cytoplasm. Nucleus.


Tissue Location

Highest expression of UBASH3A in tissues belonging to the immune system, including spleen, peripheral blood leukocytes, thymus and bone marrow.


UBASH3A / CLIP4 Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UBASH3A / CLIP4 Antibody (Internal) - Images

UBASH3A antibody (1 ug/ml) staining of Human Spleen lysate (35 ug protein/ml in RIPA buffer).



Anti-UBASH3A antibody IHC of human breast.

UBASH3A / CLIP4 Antibody (Internal) - Background

Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors, EGFR and PDGFRB,

on the cell surface. Exhibits negligible protein tyrosine phosphatase activity at neutral pH. May act as a dominant-negative regulator of UBASH3B-dependent dephosphorylation. May inhibit dynamin-dependent endocytic pathways by functionally sequestering dynamin via its SH3 domain.

UBASH3A / CLIP4 Antibody (Internal) - References

Wattenhofer M., et al. Hum. Genet. 108:140-147(2001).
Feshchenko E.A., et al. Oncogene 23:4690-4706(2004).
Hattori M., et al. Nature 405:311-319(2000).
Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Kowanetz K., et al. J. Biol. Chem. 279:32786-32795(2004).