

ALDH3A1 Antibody (C-Terminus)
Goat Polyclonal Antibody
Catalog # ALS15788**Specification**

ALDH3A1 Antibody (C-Terminus) - Product Information

Application	IHC, WB
Primary Accession	P30838
Reactivity	Human, Mouse
Host	Goat
Clonality	Polyclonal
Calculated MW	50kDa KDa

ALDH3A1 Antibody (C-Terminus) - Additional Information**Gene ID** 218**Other Names**

Aldehyde dehydrogenase, dimeric NADP-preferring, 1.2.1.5, ALDHIII, Aldehyde dehydrogenase 3, Aldehyde dehydrogenase family 3 member A1, ALDH3A1, ALDH3

Target/Specificity

Human ALDH3A1. Reported variants represent identical protein: NP_001128639.1, NP_000682.3, NP_001128640.1

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

ALDH3A1 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

ALDH3A1 Antibody (C-Terminus) - Protein Information**Name** ALDH3A1**Synonyms** ALDH3**Function**

ALDHs play a major role in the detoxification of alcohol- derived acetaldehyde (Probable). They are involved in the metabolism of corticosteroids, biogenic amines, neurotransmitters, and lipid peroxidation (Probable). Oxidizes medium and long chain aldehydes into non-toxic fatty acids (PubMed:1737758). Preferentially oxidizes aromatic aldehyde substrates (PubMed:1737758). Comprises about 50 percent of corneal epithelial soluble proteins (By similarity). May play a role in preventing corneal damage caused by ultraviolet light (By similarity).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:P47739}.


Tissue Location

High levels in stomach, esophagus and lung; low level in the liver and kidney

ALDH3A1 Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ALDH3A1 Antibody (C-Terminus) - Images

Anti-ALDH3A1 antibody IHC staining of human lung, respiratory epithelium.



ALDH3A1 antibody (0.01 ug/ml) staining of Mouse Eye lysate (35 ug protein/ml in RIPA buffer).

ALDH3A1 Antibody (C-Terminus) - Background

ALDHs play a major role in the detoxification of alcohol-derived acetaldehyde. They are involved in the metabolism of corticosteroids, biogenic amines, neurotransmitters, and lipid peroxidation. This protein preferentially oxidizes aromatic aldehyde substrates. It may play a role in the oxidation of toxic aldehydes.

ALDH3A1 Antibody (C-Terminus) - References

Schuuring E.M.D.,et al.Submitted (AUG-1991) to the EMBL/GenBank/DDBJ databases.
Hsu L.C.,et al.J. Biol. Chem. 267:3030-3037(1992).
Hsu L.C.,et al.Adv. Exp. Med. Biol. 328:141-152(1993).
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).