

MOB2 Antibody (Internal)
Goat Polyclonal Antibody
Catalog # ALS15858**Specification**

MOB2 Antibody (Internal) - Product Information

Application	IHC, WB
Primary Accession	Q70IA6
Reactivity	Human, Mouse, Rat, Hamster, Monkey, Horse, Xenopus, Bovine, Dog
Host	Goat
Clonality	Polyclonal
Calculated MW	27kDa KDa

MOB2 Antibody (Internal) - Additional Information**Gene ID** 81532**Other Names**

MOB kinase activator 2, HCCA2, Mob2 homolog, Mps one binder kinase activator-like 2, MOB2, HCCA2

Target/Specificity

Human MOB2 / HCCA2. This antibody is expected to recognize both reported isoforms (NP_001165694.1; NP_443731.2) only.

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

MOB2 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

MOB2 Antibody (Internal) - Protein Information**Name** MOB2**Synonyms** HCCA2**Function**

Stimulates the autophosphorylation and kinase activity of STK38 and STK38L.

Cellular Location


Nucleus. Cytoplasm, perinuclear region

MOB2 Antibody (Internal) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MOB2 Antibody (Internal) - Images



Anti-MOB2 antibody IHC staining of human lung, airway.



MOB2 antibody (0.02 ug/ml) staining of Jurkat lysate (35 ug protein/ml in RIPA buffer).

MOB2 Antibody (Internal) - Background

Stimulates the autophosphorylation and kinase activity of STK38 and STK38L.

MOB2 Antibody (Internal) - References

Florindo C.S.,et al.Submitted (AUG-2003) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Taylor T.D.,et al.Nature 440:497-500(2006).
Devroe E.,et al.J. Biol. Chem. 279:24444-24451(2004).
Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).

