

BRCA1 Antibody (aa1847-1863)
Rabbit Polyclonal Antibody
Catalog # ALS15962**Specification**

BRCA1 Antibody (aa1847-1863) - Product Information

Application	IHC
Primary Accession	P38398
Reactivity	Human, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	208kDa KDa

BRCA1 Antibody (aa1847-1863) - Additional Information**Gene ID** 672**Other Names**

Breast cancer type 1 susceptibility protein, 6.3.2.-, RING finger protein 53, BRCA1, RNF53

Target/Specificity

Recognizes BRCA1 also known as Breast Cancer 1, a tumor suppressor gene which is involved in the repair of damaged DNA. BRCA1 protein inhibits the growth of cells lining the milk ducts in the breast and is often used as a diagnostic marker for breast ...

Reconstitution & Storage

For long term, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

BRCA1 Antibody (aa1847-1863) is for research use only and not for use in diagnostic or therapeutic procedures.

BRCA1 Antibody (aa1847-1863) - Protein Information**Name** BRCA1**Synonyms** RNF53**Function**

E3 ubiquitin-protein ligase that specifically mediates the formation of 'Lys-6'-linked polyubiquitin chains and plays a central role in DNA repair by facilitating cellular responses to DNA damage (PubMed: 12890688, PubMed: 14976165, PubMed: 16818604, PubMed: 17525340, PubMed: 12887909, PubMed: 10500182, PubMed: 19261748).

It is unclear whether it also mediates the formation of other types of polyubiquitin chains (PubMed:12890688). The BRCA1-BARD1 heterodimer coordinates a diverse range of cellular pathways such as DNA damage repair, ubiquitination and transcriptional regulation to maintain genomic stability (PubMed:12890688, PubMed:14976165, PubMed:20351172). Regulates centrosomal microtubule nucleation (PubMed:18056443). Required for appropriate cell cycle arrests after ionizing irradiation in both the S-phase and the G2 phase of the cell cycle (PubMed:10724175, PubMed:12183412, PubMed:11836499, PubMed:19261748). Required for FANCD2 targeting to sites of DNA damage (PubMed:12887909). Inhibits lipid synthesis by binding to inactive phosphorylated ACACA and preventing its dephosphorylation (PubMed:16326698). Contributes to homologous recombination repair (HRR) via its direct interaction with PALB2, fine-tunes recombinational repair partly through its modulatory role in the PALB2-dependent loading of BRCA2-RAD51 repair machinery at DNA breaks (PubMed:19369211). Component of the BRCA1-RBBP8 complex which regulates CHEK1 activation and controls cell cycle G2/M checkpoints on DNA damage via BRCA1-mediated ubiquitination of RBBP8 (PubMed:16818604). Acts as a transcriptional activator (PubMed:20160719).

Cellular Location

Nucleus. Chromosome. Cytoplasm. Note=Localizes at sites of DNA damage at double-strand breaks (DSBs); recruitment to DNA damage sites is mediated by ABRAXAS1 and the BRCA1-A complex (PubMed:26778126) Translocated to the cytoplasm during UV-induced apoptosis (PubMed:20160719). [Isoform 5]: Cytoplasm

Tissue Location

Isoform 1 and isoform 3 are widely expressed. Isoform 3 is reduced or absent in several breast and ovarian cancer cell lines

Volume


50 µl

BRCA1 Antibody (aa1847-1863) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BRCA1 Antibody (aa1847-1863) - Images



Anti-BRCA1 antibody IHC staining of human testis.

BRCA1 Antibody (aa1847-1863) - Background

E3 ubiquitin-protein ligase that specifically mediates the formation of 'Lys-6'-linked polyubiquitin chains and plays a central role in DNA repair by facilitating cellular responses to DNA damage. It is unclear whether it also mediates the formation of other types of polyubiquitin chains. The E3 ubiquitin-protein ligase activity is required for its tumor suppressor function. The BRCA1-BARD1 heterodimer coordinates a diverse range of cellular pathways such as DNA damage repair, ubiquitination and transcriptional regulation to maintain genomic stability. Regulates centrosomal microtubule nucleation. Required for normal cell cycle progression from G2 to mitosis. Required for appropriate cell cycle arrests after ionizing irradiation in both the S-phase and the G2 phase of the cell cycle. Involved in transcriptional regulation of P21 in response to DNA damage. Required for FANCD2 targeting to sites of DNA damage. May function as a transcriptional regulator. Inhibits lipid synthesis by binding to inactive phosphorylated ACACA and preventing its dephosphorylation. Contributes to homologous recombination repair (HRR) via its direct interaction with PALB2, fine-tunes recombinational repair partly through its modulatory role in the PALB2-dependent loading of BRCA2-RAD51 repair machinery at DNA breaks. Component of the BRCA1-RBBP8 complex which regulates CHEK1 activation and controls cell cycle G2/M checkpoints on DNA damage via BRCA1-mediated ubiquitination of RBBP8.

BRCA1 Antibody (aa1847-1863) - References

Miki Y.,et al.Science 266:66-71(1994).
Smith T.M.,et al.Genome Res. 6:1029-1049(1996).
Wilson C.A.,et al.Oncogene 14:1-16(1997).
Holt J.T.,et al.Submitted (MAY-1997) to the EMBL/GenBank/DDBJ databases.
Raymond C.K.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.