

CXCR3 Antibody (C-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS16050**Specification**

CXCR3 Antibody (C-Terminus) - Product Information

Application	WB
Primary Accession	P49682
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41kDa KDa

CXCR3 Antibody (C-Terminus) - Additional Information**Gene ID** 2833**Other Names**

C-X-C chemokine receptor type 3, CXC-R3, CXCR-3, CKR-L2, G protein-coupled receptor 9, Interferon-inducible protein 10 receptor, IP-10 receptor, CD183, CXCR3, GPR9

Target/Specificity

CXCR3 antibody is human, mouse and rat reactive. At least two isoforms of CXCR3 are known to exist; this antibody will detect both isoforms.

Reconstitution & Storage

Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.

Precautions

CXCR3 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

CXCR3 Antibody (C-Terminus) - Protein Information**Name** CXCR3**Synonyms** GPR9**Function**

[Isoform 1]: Receptor for the C-X-C chemokine CXCL9, CXCL10 and CXCL11 and mediates the proliferation, survival and angiogenic activity of human mesangial cells (HMC) through a heterotrimeric G- protein signaling pathway (PubMed:12782716). Binds to CCL21. Probably promotes cell chemotaxis response. [Isoform 3]: Mediates the activity of CXCL11.

Cellular Location

[Isoform 1]: Cell membrane; Multi-pass membrane protein

Tissue Location


Isoform 1 and isoform 2 are mainly expressed in heart, kidney, liver and skeletal muscle. Isoform 1 is also expressed in placenta. Isoform 2 is expressed in endothelial cells. Expressed in T-cells (at protein level).

CXCR3 Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CXCR3 Antibody (C-Terminus) - Images



Western blot analysis of CXCR3 in HepG2 lysate with CXCR3 antibody at (A) 1 and (B) 2 ug/ml.

CXCR3 Antibody (C-Terminus) - Background

Isoform 1: Receptor for the C-X-C chemokine CXCL9, CXCL10 and CXCL11 and mediates the proliferation, survival and angiogenic activity of human mesangial cells (HMC) through a heterotrimeric G-protein signaling pathway (PubMed:12782716). Binds to CCL21. Probably promotes cell chemotaxis response. Isoform 3: Mediates the activity of CXCL11.

CXCR3 Antibody (C-Terminus) - References

Loetscher M., et al. J. Exp. Med. 184:963-969(1996).
Lasagni L., et al. J. Exp. Med. 197:1537-1549(2003).
Petrai I., et al. Int. J. Biochem. Cell Biol. 40:1764-1774(2008).
Gutierrez J., et al. Submitted (SEP-1996) to the EMBL/GenBank/DDBJ databases.
Warren C.N., et al. Submitted (FEB-2003) to the EMBL/GenBank/DDBJ databases.