

CAV1 / Caveolin 1 Antibody (aa129-178) Rabbit Polyclonal Antibody Catalog # ALS16134

Specification

CAV1 / Caveolin 1 Antibody (aa129-178) - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW IHC, IF, WB <u>003135</u> Human, Mouse, Rat Rabbit Polyclonal 20kDa KDa

CAV1 / Caveolin 1 Antibody (aa129-178) - Additional Information

Gene ID 857

Other Names Caveolin-1, CAV1, CAV

Target/Specificity Caveolin-1 Antibody detects endogenous levels of total Caveolin-1 protein.

Reconstitution & Storage Store at -20°C for up to one year.

Precautions CAV1 / Caveolin 1 Antibody (aa129-178) is for research use only and not for use in diagnostic or therapeutic procedures.

CAV1 / Caveolin 1 Antibody (aa129-178) - Protein Information

Name CAV1

Synonyms CAV

Function

May act as a scaffolding protein within caveolar membranes (PubMed:11751885). Forms a stable heterooligomeric complex with CAV2 that targets to lipid rafts and drives caveolae formation. Mediates the recruitment of CAVIN proteins (CAVIN1/2/3/4) to the caveolae (PubMed:19262564). Interacts directly with G-protein alpha subunits and can functionally regulate their activity (By similarity). Involved in the costimulatory signal essential for T-cell receptor (TCR)-mediated T-cell activation. Its binding to DPP4 induces T-cell proliferation and NF-kappa-B activation in a T-cell receptor/CD3-dependent manner (PubMed:17287217). Recruits CTNNB1 to caveolar membranes and may regulate CTNNB1-mediated signaling through the Wnt pathway (By similarity). Negatively regulates



TGFB1-mediated activation of SMAD2/3 by mediating the internalization of TGFBR1 from membrane rafts leading to its subsequent degradation (PubMed:25893292). Binds 20(S)-hydroxycholesterol (20(S)-OHC) (By similarity).

Cellular Location

Golgi apparatus membrane; Peripheral membrane protein. Cell membrane; Peripheral membrane protein. Membrane, caveola; Peripheral membrane protein. Membrane raft. Golgi apparatus, trans-Golgi network {ECO:0000250|UniProtKB:P33724} Note=Colocalized with DPP4 in membrane rafts. Potential hairpin-like structure in the membrane. Membrane protein of caveolae

Tissue Location

Skeletal muscle, liver, stomach, lung, kidney and heart (at protein level). Expressed in the brain


Volume 50 μl

CAV1 / Caveolin 1 Antibody (aa129-178) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

CAV1 / Caveolin 1 Antibody (aa129-178) - Images




Anti-CAV1 / Caveolin 1 antibody IHC staining of human liver.



Immunofluorescence of HUVEC cells, using Caveolin-1 Antibody.





Western blot of extracts from HUVEC cells, using Caveolin-1 Antibody.

CAV1 / Caveolin 1 Antibody (aa129-178) - Background

May act as a scaffolding protein within caveolar membranes. Interacts directly with G-protein alpha subunits and can functionally regulate their activity (By similarity). Involved in the costimulatory signal essential for T-cell receptor (TCR)- mediated T-cell activation. Its binding to DPP4 induces T-cell proliferation and NF-kappa-B activation in a T-cell receptor/CD3- dependent manner. Recruits CTNNB1 to caveolar membranes and may regulate CTNNB1-mediated signaling through the Wnt pathway.

CAV1 / Caveolin 1 Antibody (aa129-178) - References

Glenney J.R. Jr., et al.FEBS Lett. 314:45-48(1992). Hurlstone A.F., et al.Oncogene 18:1881-1890(1999). Engelman J.A., et al.FEBS Lett. 448:221-230(1999). Kalnine N., et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases. Vainonen J.P., et al.Biochem. Biophys. Res. Commun. 320:480-486(2004).