10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CREB3 / LZIP Antibody (aa76-87)

Goat Polyclonal Antibody
Catalog # ALS16145

Specification

CREB3 / LZIP Antibody (aa76-87) - Product Information

Application	**WB**
Primary Accession	O43889
Reactivity	**Human**
Host	**Goat**
Clonality	**Polyclonal**
Calculated MW	**44kDa KDa**

CREB3 / LZIP Antibody (aa76-87) - Additional Information

Gene ID 10488

Other Names
Cyclic AMP-responsive element-binding protein 3, CREB-3, cAMP-responsive element-binding protein 3, Leucine zipper protein, Luman, Transcription factor LZIP-alpha, Processed cyclic AMP-responsive element-binding protein 3, N-terminal Luman, Transcriptionally active form, CREB3, LZIP

Target/Specificity
Human CREB3 / LZIP

Reconstitution & Storage
Store at -20°C. Minimize freezing and thawing.

Precautions
CREB3 / LZIP Antibody (aa76-87) is for research use only and not for use in diagnostic or therapeutic procedures.

CREB3 / LZIP Antibody (aa76-87) - Protein Information

Name CREB3

Synonyms LZIP

Function
Endoplasmic reticulum (ER)-bound sequence-specific transcription factor that directly binds DNA and activates transcription (PubMed:9271389, PubMed:19779205, PubMed:10984507, PubMed:15845366, PubMed:16940180). Plays a role in the unfolded protein response (UPR), promoting cell survival versus ER stress-induced apoptotic cell death (PubMed:<a

href="http://www.uniprot.org/citations/15845366" target="_blank">15845366, PubMed:16940180). Also involved in cell proliferation, migration and differentiation, tumor suppression and inflammatory gene expression. Acts as a positive regulator of LKN- 1/CCL15-induced chemotaxis signaling of leukocyte cell migration (PubMed:19779205, PubMed:15001559, PubMed:17296613). Associates with chromatin to the HERPUD1 promoter (PubMed:16940180). Also induces transcriptional activation of chemokine receptors (PubMed:18587271, PubMed:17296613).

Cellular Location
[Isoform 1]: Endoplasmic reticulum membrane; Single-pass type II membrane protein {ECO:0000255, ECO:0000269|PubMed:12138176}. Golgi apparatus. Note=Colocalizes with HCFC1 in neuronal cell bodies of the trigeminal ganglia (PubMed:10623756). Colocalizes with DCSTAMP in the ER membrane of immature dendritic cell (DC) (PubMed:20546900). Colocalizes with CANX, CCR1, HCFC1 in the ER membrane (PubMed:10623756). [Isoform 2]: Nucleus. Cytoplasm Note=Predominantly in the nucleus (PubMed:19779205). Not associated with membranes (PubMed:19779205).

Tissue Location
Ubiquitously expressed (PubMed:9271389, PubMed:19779205). Expressed in dendritic cells (DC). Weakly expressed in monocytes (at protein level) (PubMed:20546900)

CREB3 / LZIP Antibody (aa76-87) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CREB3 / LZIP Antibody (aa76-87) - Images



10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CREB3 antibody (0.3 ug/ml) staining of HepG2 lysate (35 ug protein in RIPA buffer).

CREB3 / LZIP Antibody (aa76-87) - Background

Endoplasmic reticulum (ER)-bound transcription factor that plays a role in the unfolded protein response (UPR). Involved in cell proliferation and migration, tumor suppression and inflammatory gene expression. Plays also a role in the human immunodeficiency virus type 1 (HIV-1) virus protein expression and in the herpes simplex virus-1 (HSV-1) latent infection and reactivation from latency. Isoform 2 plays a role in the unfolded protein response (UPR). Isoform 2 acts as a positive regulator of LKN-1/CCL15-induced chemotaxis signaling of leukocyte cell migration. Isoform 2 may play a role as a cellular tumor suppressor that is targeted by the hepatitis C virus (HSV) core protein. Isoform 2 represses the VP16-mediated transactivation of immediate early genes of the HSV-1 virus by sequestring host cell factor-1 HCFC1 in the ER membrane of sensory neurons, thereby preventing the initiation of the replicative cascade leading to latent infection. Isoform 3 functions as a negative transcriptional regulator in ligand-induced transcriptional activation of the glucocorticoid receptor NR3C1 by recruiting and activating histone deacetylases (HDAC1, HDAC2 and HDAC6). Isoform 3 decreases the acetylation level of histone H4. Isoform 3 does not promote the chemotactic activity of leukocyte cells.

CREB3 / LZIP Antibody (aa76-87) - References

Lu R.,et al.Mol. Cell. Biol. 17:5117-5126(1997).
Freiman R.N.,et al.Genes Dev. 11:3122-3127(1997).
Jin D.-Y.,et al.EMBO J. 19:729-740(2000).
Kang H.,et al.Mol. Endocrinol. 23:1746-1757(2009).
Hayashi M.,et al.Submitted (FEB-1997) to the EMBL/GenBank/DDBJ databases.