

CENPU / MLF1IP Antibody (C-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS16165**Specification****CENPU / MLF1IP Antibody (C-Terminus) - Product Information**

Application	IHC
Primary Accession	Q71F23
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48kDa KDa

CENPU / MLF1IP Antibody (C-Terminus) - Additional Information**Gene ID** 79682**Other Names**

Centromere protein U, CENP-U, Centromere protein of 50 kDa, CENP-50, Interphase centromere complex protein 24, KSHV latent nuclear antigen-interacting protein 1, MLF1-interacting protein, Polo-box-interacting protein 1, CENPU, ICEN24, KLIP1, MLF1IP, PBIP1

Target/Specificity

A BLAST analysis was used to suggest cross-reactivity with MLF1IP protein from human, chimpanzee (98%), horse (72%), bovine (69%), dog (65%), mouse (56%), rat (55%) and chicken (47%) sources based on homology with the immunizing sequence. Reactivity ...

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.

Precautions

CENPU / MLF1IP Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

CENPU / MLF1IP Antibody (C-Terminus) - Protein Information**Name** CENPU**Synonyms** ICEN24, KLIP1, MLF1IP, PBIP1**Function**

Component of the CENPA-NAC (nucleosome-associated) complex, a complex that plays a central role in assembly of kinetochore proteins, mitotic progression and chromosome segregation. The CENPA-NAC complex recruits the CENPA-CAD (nucleosome distal) complex and may be involved in incorporation of newly synthesized CENPA into centromeres. Plays an important role in the correct PLK1 localization to the mitotic kinetochores. A scaffold protein responsible for the initial recruitment and maintenance of the kinetochore PLK1 population until its degradation. Involved in transcriptional repression.

Cellular Location

Cytoplasm. Nucleus. Chromosome, centromere, kinetochore. Note=Localizes in the kinetochore domain of centromeres Colocalizes with PLK1 at the interzone between the inner and the outer kinetochore plates


Tissue Location

Expressed at high levels in the testis, fetal liver, thymus, bone marrow and at lower levels in the lymph nodes, placenta, colon and spleen. Present in all cell lines examined, including B-cells, T-cells, epithelial cells and fibroblast cells Expressed at high levels in glioblastoma cell lines

CENPU / MLF1IP Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CENPU / MLF1IP Antibody (C-Terminus) - Images

Anti-CENPU / MLF1IP antibody IHC staining of human testis.

CENPU / MLF1IP Antibody (C-Terminus) - Background

Component of the CENPA-NAC (nucleosome-associated) complex, a complex that plays a central role in assembly of kinetochore proteins, mitotic progression and chromosome segregation. The CENPA-NAC complex recruits the CENPA-CAD (nucleosome distal) complex and may be involved in incorporation of newly synthesized CENPA into centromeres. Plays an important role in the correct PLK1 localization to the mitotic kinetochores. A scaffold protein responsible for the initial recruitment and maintenance of the kinetochore PLK1 population until its degradation. Involved in transcriptional repression.

CENPU / MLF1IP Antibody (C-Terminus) - References

Pan H.-Y., et al. J. Virol. 77:9758-9768(2003).
Hanissian S.H., et al. Oncogene 23:3700-3707(2004).

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.

Hanissian S.H.,et al.Submitted (AUG-2006) to the EMBL/GenBank/DDBJ databases.