

**HTR2C / 5-HT2C Receptor Antibody (aa400-450)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS16334****Specification**

---

**HTR2C / 5-HT2C Receptor Antibody (aa400-450) - Product Information**

Application	IHC
Primary Accession	<a href="#">P28335</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	52kDa KDa

**HTR2C / 5-HT2C Receptor Antibody (aa400-450) - Additional Information****Gene ID** 3358**Other Names**

5-hydroxytryptamine receptor 2C, 5-HT-2C, 5-HT2C, 5-HTR2C, 5-hydroxytryptamine receptor 1C, 5-HT-1C, 5-HT1C, Serotonin receptor 2C, HTR2C, HTR1C

**Target/Specificity**

Human HTR2C / 5-HT2C

**Reconstitution & Storage**

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

**Precautions**

HTR2C / 5-HT2C Receptor Antibody (aa400-450) is for research use only and not for use in diagnostic or therapeutic procedures.

**HTR2C / 5-HT2C Receptor Antibody (aa400-450) - Protein Information****Name** HTR2C ([HGNC:5295](#))**Synonyms** HTR1C**Function**

G-protein coupled receptor for 5-hydroxytryptamine (serotonin). Also functions as a receptor for various drugs and psychoactive substances, including ergot alkaloid derivatives, 1-2,5,-dimethoxy-4-iodophenyl-2-aminopropane (DOI) and lysergic acid diethylamide (LSD). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors. Beta-arrestin family members inhibit signaling via G proteins and mediate activation of alternative signaling pathways. Signaling activates a phosphatidylinositol-calcium second messenger system that modulates the activity of phosphatidylinositol 3-kinase and down-stream signaling cascades and promotes the release of Ca(2+) ions from intracellular stores. Regulates neuronal activity via the activation of short transient receptor potential calcium channels in the brain, and thereby modulates the

activation of pro-opiomelanocortin neurons and the release of CRH that then regulates the release of corticosterone. Plays a role in the regulation of appetite and eating behavior, responses to anxiogenic stimuli and stress. Plays a role in insulin sensitivity and glucose homeostasis.

**Cellular Location**

Cell membrane; Multi-pass membrane protein


**Tissue Location**

Detected in brain..

**HTR2C / 5-HT2C Receptor Antibody (aa400-450) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**HTR2C / 5-HT2C Receptor Antibody (aa400-450) - Images**

Anti-HTR2C / 5-HT2C Receptor antibody IHC staining of human brain, cerebellum.

**HTR2C / 5-HT2C Receptor Antibody (aa400-450) - Background**

G-protein coupled receptor for 5-hydroxytryptamine (serotonin). Also functions as a receptor for various drugs and psychoactive substances, including ergot alkaloid derivatives, 1-2,5,-dimethoxy-4-iodophenyl-2-aminopropane (DOI) and lysergic acid diethylamide (LSD). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors. Beta-arrestin family members inhibit signaling via G proteins and mediate activation of alternative signaling pathways. Signaling activates a phosphatidylinositol-calcium second messenger system that modulates the activity of phosphatidylinositol 3-kinase and down-stream signaling cascades and promotes the release of  $\text{Ca}^{2+}$  ions from intracellular stores. Regulates neuronal activity via the activation of short transient receptor potential calcium channels in the brain, and thereby modulates the activation of pro-opiomelanocortin neurons and the release of CRH that then regulates the release of corticosterone. Plays a role in the regulation of appetite and eating behavior, responses to

angiogenic stimuli and stress. Plays a role in insulin sensitivity and glucose homeostasis.

#### **HTR2C / 5-HT2C Receptor Antibody (aa400-450) - References**

- Saltzman A.G., et al. Biochem. Biophys. Res. Commun. 181:1469-1478(1991).  
Stam N.J., et al. Eur. J. Pharmacol. 269:339-348(1994).  
Xie E., et al. Genomics 35:551-561(1996).  
Niswender C.M., et al. Ann. N. Y. Acad. Sci. 861:38-48(1998).  
Puhl H.L. III, et al. Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases.