10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

E2F5 Antibody (aa93-142)

Rabbit Polyclonal Antibody

Catalog # ALS16341

Specification

E2F5 Antibody (aa93-142) - Product Information

Application	**WB**
Primary Accession	Q15329
Reactivity	**Human, Mouse, Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**38kDa KDa**

E2F5 Antibody (aa93-142) - Additional Information

Gene ID 1875

Other Names
Transcription factor E2F5, E2F-5, E2F5

Target/Specificity
E2F-5 antibody detects endogenous levels of E2F-5.

Reconstitution & Storage
Store at -20°C.

Precautions
E2F5 Antibody (aa93-142) is for research use only and not for use in diagnostic or therapeutic procedures.

E2F5 Antibody (aa93-142) - Protein Information

Name E2F5

Function
Transcriptional activator that binds to E2F sites, these sites are present in the promoter of many genes whose products are involved in cell proliferation. May mediate growth factor-initiated signal transduction. It is likely involved in the early responses of resting cells to growth factor stimulation. Specifically required for multiciliate cell differentiation: together with MCIDAS and E2F5, binds and activate genes required for centriole biogenesis.

Cellular Location
Nucleus.

Volume
50 µl


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

E2F5 Antibody (aa93-142) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

E2F5 Antibody (aa93-142) - Images



Western blot of E2F-5 antibody in extracts from HeLa cells.

E2F5 Antibody (aa93-142) - Background

Transcriptional activator that binds to E2F sites, these sites are present in the promoter of many genes whose products are involved in cell proliferation. May mediate growth factor- initiated signal transduction. It is likely involved in the early responses of resting cells to growth factor stimulation. Specifically required for multiciliate cell differentiation: together with MCIDAS and E2F5, binds and activate genes required for centriole biogenesis.

E2F5 Antibody (aa93-142) - References

Itoh A.,et al.Cell. Mol. Biol. Res. 41:147-154(1995).
Hijmans E.M.,et al.Mol. Cell. Biol. 15:3082-3089(1995).
Sardet C.,et al.Proc. Natl. Acad. Sci. U.S.A. 92:2403-2407(1995).
Vaishnav Y.N.,et al.Biochem. Biophys. Res. Commun. 242:586-592(1998).
Li W.B.,et al.Submitted (APR-2003) to the EMBL/GenBank/DDBJ databases.